



Hudsonia
a nonprofit institute

Field Station: PO Box 5000, Annandale, NY 12504-5000
Phone: (845) 758-7053
Fax: (845) 758-7033

Red Hook: PO Box 66, Red Hook, NY 12571
Phone: (845) 758-0600
Fax: (845) 758-0607

www.hudsonia.org

Experimental Management of Japanese Knotweed
(*Fallopia japonica*)
on the Batavia Kill, Greene County, New York



by

Tanessa Hartwig and Erik Kiviat
Hudsonia Ltd.
Annandale, NY

Report to

Greene County Soil and Water Conservation District
and
New York City Department of Environmental Protection

December 2009

INTRODUCTION

Hudsonia Ltd. was asked by the Greene County Soil and Water Conservation District (GCSWCD) and the New York City Department of Environmental Protection (NYCDEP) to test management techniques that could be used by landowners to effectively control Japanese knotweed (Polygonaceae: *Fallopia japonica* [*Polygonum cuspidatum*], hereinafter “knotweed”) on a local scale. Knotweed stands are widespread, and in many places extensive, on the banks and floodplain of the Batavia Kill in Greene County, New York. The Batavia Kill flows into Schoharie Creek which feeds Schoharie Reservoir, part of the New York City (NYC) water supply system. There is a high level of concern about the potential impacts of knotweed on water quality in streams in the NYC water supply watershed (Talmage and Kiviat 2004). Our study focused on three management techniques – partial replacement of knotweed with woody plantings, herbicide injection into uncut stems, and frequent cutting – at a study site on the banks of the Batavia Kill. This report presents the results of these studies and recommendations for management of knotweed. The GCSWCD and NYCDEP intend to use this study to develop guidance for property owners who wish to manage knotweed on their own lands.

Hudsonia Ltd. is an independent, non-advocacy, non-profit institute for research and education in the environmental sciences. We conduct studies to assess biological resources and make recommendations for ecologically sound land management. Hudsonia does not support or oppose land use projects; rather we provide scientific information, analyses, and recommendations impartially for use by parties involved in environmental planning and environmental management.

Metric units of measurement are used in this report. English equivalents are:

- 1 mm (millimeter) = 0.04 inch
- 1 cm (centimeter) = 0.39 inch
- 1 m (meter) = 3.28 feet
- 1 mL (milliliter) = 0.03 ounces

JAPANESE KNOTWEED

Knotweed was probably introduced to North America in the late 1800s (Seiger 1997, Child and Wade 2000), and is now widespread and invasive especially in riparian environments and on human-disturbed soils where it can form dense stands with a depauperate (species-poor) ground flora. Knotweed is native to Japan, northern China, Korea, and Taiwan (Child & Wade 2000). It is usually considered herbaceous and has annual aerial stems that die back to near ground level in autumn, and long-lived perennial root systems. Knotweed grows to 1-3 or more meters high. It has broadly egg-shaped, alternate leaves and greenish-white flowers arranged in clusters in the leaf axils.

Knotweed appears to root more shallowly than native riparian trees and shrubs, although more deeply than the local grasses. Streambanks dominated by knotweed, therefore, may be less stable and more prone to slumping and erosion than banks with trees or shrubs (Rene VanSchaack, personal communication). This is believed to increase suspended sediment loads and turbidity, and thus degrade water quality in the Batavia Kill. Because more turbid water is more expensive to treat and more likely to harbor pathogens,

knotweed is a concern for the NYC water supply system. Populations of knotweed have also been reported on other streams of the Catskill Mountain component of the NYC water supply watershed, and these populations might also affect water quality. Rigorous scientific study of the relationship of knotweed to bank stability and water quality has not been conducted, as far as we know.

In addition to the potential effects of knotweed on water quality, the plant could also affect fisheries. This could occur via alteration of the detritus (dead plant matter) based food web that supports trout and other stream fishes, as well as via interference with human access to stream banks for angling. Because riparian zones are a critical component of the environment for many other animal and plant species, physical, chemical, and biological changes to the riparian zones caused by a large-scale plant invasion could also change habitat functions, including habitat for plants, and food, shelter, and nesting sites for animals. Geographically and taxonomically wide ranging natural history study in the northeastern states and adjacent Canada indicate a complex biota of native and introduced species of animals, plants, and fungi associated with stands of knotweed (Kiviat, unpublished data). However, methodologically sound quantitative research on knotweed effects on the fitness and population dynamics of a broad range of native species remains to be conducted.

STUDY AREA

Figure 1 illustrates the study area. Four plot types (control, woody planting, herbicide injection, and frequent cutting) were located on the south bank of the Batavia Kill in the Town of Ashland in the northern Catskill Mountains in Greene County, New York, ca. 60 km southwest of Albany, New York (U.S. Geological Survey, 1945 Ashland 7.5 minute topographic map sheet). The study site was ca. 400 m west (downstream) of County Route 17. At this location, the Batavia Kill is classified as a class C trout stream (NYSDEC 2009). Soils are Barbour loam, a very deep, well-drained soil that formed in alluvial deposits (Day 2003). Soils are underlain by the Oneonta shale, sandstone, and conglomerate formation. Topography at the site is nearly level except where the bank slopes steeply to the creek. Land uses surrounding the study site were primarily agricultural; fields next to the plots were planted with corn during the study.

METHODS

Plot location

Nine 5 x 5 m plots were permanently marked on the floodplain of the Batavia Kill. (Ten additional plots were originally established on a nearby site; however, the additional plots were compromised due to the activities of a landowner.) Plots were randomly spaced 3-7 m apart to avoid potential impacts from neighboring treatment plots and to allow equipment access between plots. The four treatments (control, woody planting, herbicide injection, and frequent cutting) were assigned randomly. Plots were at least 1 m inside the stand edge, to avoid the weedy grasses and forbs associated with the stand edges. All plots were in stands in which knotweed was highly dominant, with the streamside plot edge 0-16 m (\bar{x} = 10 m) from the top of the stream bank. We did not locate plots beneath substantial crown cover of tall woody plants. Knotweed is dioecous; in our plots all shoots examined were female.

Pre-treatment surveys

The vegetation on the plots was sampled 21 August to 5 September 2003, and the soils were sampled 16-26 September 2003, before treatments were initiated. For each whole plot, we visually estimated percent knotweed canopy cover, percent litter cover (all species), percent bryophyte cover on the soil, and percent insect damage to knotweed leaves. We identified all other vascular plants and visually estimated the percent cover of each species and its maximum height (cm). Because the garlic-mustard (*Alliaria petiolata*) senesced during the period of sampling, cover was estimated as if the associated species were in full summer condition. We recorded the maximum height of the knotweed canopy over three of the plot corners (excluding the first corner established for each plot) with a telescoping tree measuring pole. We measured basal diameter (10 cm above the soil, in mm) of all knotweed stems in 1 x 1 m subplots within the same three corners of the plot using a diameter tape.

Soil characteristics were observed in a soil pit in the middle of the western boundary of each 5 x 5 m plot. We avoided digging inside the plots because of their intended use for management experiments. Pits were dug with a shovel to a depth of ca. 1 m, and deeper characteristics were examined using a Dutch auger to the lower boundary of the C horizon when possible. For each soil horizon, observations on horizon thickness, color, pH, root distribution, and percent rock fragments were recorded (see Shoeneberger et al. 2002). Color was determined by comparison with a Munsell soil color chart (Kollmorgen Corp., Newburgh, New York). pH was estimated with the Cornell pH test kit with an accuracy of 0.2 to 0.4 pH unit.

GCSWCD and NYCDEP staff marked, trenched, and treated the plots. Larry Day (Delaware County Soil and Water Conservation District) performed the soil study.

Treatment techniques

Site preparation: To prevent the subsurface incursion of knotweed from outside the plots, trenches 0.6 m wide by 0.6 m deep were excavated adjoining the periphery of each plot. Each trench was then lined with 2 layers of silt fence material (filter fabric) and backfilled with clean soil when available, or with excavated material cleansed of knotweed clumps. Plots were prepared 21-23 September 2004.

Woody planting plots: Five woody plants were placed in each plot – a silver maple (*Acer saccharinum*), a box elder (*Acer negundo*), a red maple (*Acer rubrum*), a green ash (*Fraxinus pennsylvanica*), and a black cherry (*Prunus serotina*). Knotweed was grubbed from a 1 m diameter circle and the soil was excavated deep enough to accommodate the root ball of the planting stock. Each tree was then placed within a circle, and the area backfilled with knotweed-free soil. Two layers of 1 m diameter porous weed mat were placed around the planted tree. Each species was planted in the same location (the four corners and the center) on each plot. Plantings took place 2-4 May 2005. Planted trees were at least 1.2 m tall. Three plots received this treatment.

Frequent cutting plots: In 2005 and 2006, knotweed stems were cut to 20-25 cm above the ground using a metal-bladed Stihl SS250 Weed Eater approximately every other week (8 and 7 times, respectively) during the growing season (from late May to late August). In 2007, plots were cut every other week (3 times) from late May to late June. Cut

knotweed stalks were removed from the site, dried, and burned. Two plots received this treatment.

Herbicide injection plots: In mid-August 2005, an opening was cut just below the first or second node above ground-level of all stems greater than 10 mm diameter. The opening was cut using a Japanese knotweed probe (JK International, www.jkinjectiontools.com). Five mL of the herbicide glyphosate (Aqua-Master, www.monsanto.com) was then injected into the opening. Methods were adopted from Crockett et al. 2004. Two plots received this treatment. Herbicide injections were repeated in mid-August 2008.

Post-treatment surveys

All vegetation variables sampled pre-treatment (2003) were sampled post-treatment 21 August 2006, and 26-28 August and 7 September 2009. Bryophyte cover was not estimated in 2006 but was in 2009. In 2006 we collected bryophytes from frequently cut plots and in 2009 we collected bryophytes from all plots where they occurred, for later determination. In 2009 litter depths were also measured, to the nearest cm.

For consistency, Erik Kiviat and Jennifer Grieser sampled the vegetation in all three years. Vascular plant scientific names used in the report are based on the New York Flora Atlas (www.newyork.plantatlas.usf.edu). Vascular plants that could not be identified with confidence in the field were collected and the specimens identified by James (Spider) Barbour of Hudsonia. Mosses were identified by Sue Williams (Dalton, Massachusetts).

Data analysis

To compare variables by treatment and year, we created box plots. Because we only had two data points for control, frequent cutting, and herbicide injection treatments, and three points for woody planting treatments, we assigned the middle point as the median and the boxes as the minimum and maximum for each plot type. In the text, we present means and 95% confidence intervals (CI) for selected variables. Aboveground biomass was estimated from basal shoot diameters, using a regression equation derived from live basal diameter and shoot dry weight of approximately 100 untreated stems representing a wide range of size collected opportunistically from the Batavia Kill (Kiviat et al. 2004). To determine the importance of associated native and introduced plants, we multiplied percent cover by maximum height for each species and then summed the products to create indices of native and introduced plant vigor. For all calculations, < 1% cover was arbitrarily considered 0.1%. Because the differences from 2003 to 2006 and 2009 were dramatic, we believe our approach to data analysis is appropriate.

RESULTS

Pre-treatment (2003), knotweed in all plots was tall (\bar{x} = 2.9 m, 95% CI = 2.6-3.1 m) with a high percent canopy cover (\bar{x} = 85%, 95% CI = 80-90%). Average basal area was 3571 mm²/m² (95% CI = 1773-5368 mm²/m²), and average aboveground biomass was 866 g/m² (95% CI = 327-1405 g/m²). Knotweed basal stem diameter averaged 16 mm, and ranged from 11 mm to 23 mm. Litter cover was high (\bar{x} = 79%; 95% CI = 70-87%) and bryophyte cover was virtually absent; only one species was found in one plot. Average leaf tissue loss to grazing insects was minimal (\bar{x} = 1%), though maximum leaf tissue loss was high in a few instances (range = 15-60%). The number of associated

vascular plant species in each plot was low ($\bar{x} = 4$; range = 1-10; Table 1). Other introduced plants made up most of the associated species cover (native to introduced index ratio = 1:26).

Post-treatment, in the **woody planting plots**, by 2006 all woody plantings appeared dead apart from one unthrifty green ash. By 2009, all woody plantings were dead except for one 3 m silver maple. Knotweed cover was consistently high throughout the study (Figure 2), and knotweed stem density, basal area, and biomass also remained consistent (Figures 3-5). The number of associated species remained consistently low, as did the vigor of both native and introduced plants (Figures 6-8). Mean canopy height decreased slightly in woody planting plots in 2009 (Figure 9). Percent litter cover increased slightly during the course of the study (Figure 10). Results were similar in the **control** plots, though associated introduced species had somewhat more vigor in 2003 (Figures 2-10). Leaf tissue loss was low in all treatment plots throughout the study (Figure 11).

In **herbicide injection plots**, knotweed cover decreased in 2006 (post-treatment) and remained low in 2009. Stem count, canopy height, basal area, and biomass all decreased dramatically in 2006; by 2009, the knotweed had recovered, but not appreciably (Figures 3-5, 9). Litter cover decreased post-treatment, but not substantially (Figure 10). The number of associated species increased dramatically post-treatment, and remained high in 2009 (Figure 6; Tables 2 and 3). By 2009, native plant vigor was high compared with other treatments (Figure 7). Introduced plant vigor decreased immediately post-treatment, but increased to greater than pre-treatment levels by 2009 (Figure 8).

In **frequent cutting plots**, knotweed and litter cover were greatly reduced in 2006 (during treatment; Figure 2). However, by 2009 (two years post-treatment), knotweed and litter cover were almost as high as pre-treatment cover (Figures 2 and 10). Canopy heights were also greater in 2009, though they had not recovered to pre-treatment heights (Figure 9). However, basal area and biomass both remained low post-treatment (Figures 4 and 5). Stem counts increased during treatment, but then dropped to pre-treatment counts (Figure 3). The number of associated species steadily increased from pre-treatment levels during the study (Figure 6; Tables 2 and 3). Species richness was higher in the cutting plots compared to the control and herbicide plots pre-treatment (Table 1) but we feel the great increase in species numbers post-treatment was clearly due to treatment. Neither introduced nor native plants had increased vigor; in fact, the vigor of introduced plants decreased from pre-treatment levels.

On 13 November 2007, five trees were planted in each frequent cutting plot. Species planted were box elder, red maple, black cherry, and sugar maple (*Acer saccharum*). The intention was to replant the woody planting plots with faster growing and larger trees using stock grown from the “root production method” (RPM) (RPM Ecosystems, www.rpmecosystems.com); however the landscapers could not locate the woody planting plots and planted them in the cut plots instead. Several of these plantings survived in 2009; a 3 m box elder, a 0.5 m red maple, and a 1.4 m red maple in one plot and a 1.8 m box elder and a 2.4 m box elder in the second plot.

In 2009, litter depths were lower in herbicide injection and frequent cutting plots than in control and woody planting plots (Figure 12). By 2009, mosses in the cutting and

herbicide injection plots were quite diverse, particularly in the herbicide plots (Table 4). Although we did not record percent cover of bryophytes in 2006, it was clearly < 1%; however species richness of mosses was much greater than in 2003.

Species duration did not show any consistent patterns during the study (Table 5). Biennials tended to increase in herbicide injection and frequent cuttings plots, and perennials increased rather strongly in both treatment types. Woody species and annuals did not exhibit consistent patterns. However, with the small number of plots, these patterns are not strong enough to indicate relationships between duration and treatment type or duration and length of time post-treatment.

Some of the plant species that appeared in treated plots (Tables 2-3) were ecologically interesting. One species, *Eupatorium altissimum*, has previously not been found north of southern Ulster County (James Barbour, personal communication); this may indicate that the clearings created by treatment in the knotweed stands have warm microclimates. There were several wetland species (e.g., *Mimulus ringens*, *Polygonum punctatum*, *Polygonum sagittatum*, *Ludwigia palustris*) that may reflect low spots on the floodplain combined with a very wet growing season in 2009.

We found few relationships of soil characteristics to knotweed characteristics (Kiviat et al. 2004). This suggests that the floodplain soils of the study area were not limiting to the development of knotweed stands.

DISCUSSION AND RECOMMENDATIONS

In collaboration with GCSWCD and NYCDEP, we performed experimental management of Japanese knotweed on 5 x 5 meter plots using four different treatments: control, woody planting, herbicide injection, and frequent cutting. The woody planting treatment constituted five 1 m diameter circles grubbed free of knotweed in each plot with nursery stock of native trees planted one per circle. Most of the plantings died; however, taller (> 3 m instead of 1 m) or faster growing stock may have a greater chance of success.

The other two treatments consisted of herbicide injection into live stems 10 mm or greater in basal diameter and cutting several times during the growing season with removal of cut material. These treatments were successful at creating openings in the knotweed stands that were quickly colonized by diverse native and introduced herbaceous species and native mosses. Dense knotweed cover and litter apparently excludes most bryophytes and vascular plants. Once the knotweed is removed, vascular plants increase and mosses flourish on soil and decaying knotweed root crowns. Some live knotweed material persisted despite multiple years of treatment. As with most invasive plant management, treated areas need to be monitored and maintained in the long term. If treatments had continued, the vigor of remaining knotweed plants would probably have continued to decline with death resulting eventually.

Based on our results, both herbicide injection and frequent cutting have the potential to successfully manage knotweed stands on a small scale. These treatments may also have the potential to contain larger areas if used around the edges of the patches. Cutting, however, may require more maintenance or a longer treatment period than herbicide

injection; several measures of knotweed vigor (cover, stem count) indicated a return to near pre-treatment levels in the cut plots by 2009 (following cutting during 2005-June 2007). Others (biomass and basal area), however, indicated continued reduced vigor. If the cutting were continued for several more years, or if cutting were combined with other mechanical methods (e.g. shading; Seiger 1991), knotweed vigor in the cut plots may be reduced to levels equivalent to those in the herbicide plots. A greenhouse study demonstrated that cutting more than once had a significant impact on the belowground biomass of knotweed, but cutting alone did not eliminate the knotweed (Seiger & Merchant 1997). The Vermont Chapter of the Nature Conservancy reduced the height of knotweed shoots by cutting or hand-pulling knotweed four times a year for four years (Catey Ritchie, personal communication, 2002-2003). In the United Kingdom, Baker (1988) greatly reduced knotweed by cutting semimonthly during the growing season for two years.

In herbicide plots, most measures of knotweed vigor remained low, although a slight increase in certain measures (e.g., canopy height, basal area) in 2009 suggests the need for continued maintenance in these plots as well. Another study of glyphosate injection into knotweed stems also determined that complete knotweed elimination may require numerous applications over several years (Hagen and Dunwiddie 2008). Hagen and Dunwiddie (2008) also noted that stems smaller than 2 cm diameter could not receive a precise application; therefore, as knotweed stems become smaller in subsequent application seasons, the proportion of stems that can be accurately treated decreases.

In the woody planting plots, the planted trees did not survive and therefore were not successful in shading knotweed stems. We believe the plants did not thrive because they were too small to compete with the knotweed. Planting larger or faster growing trees and shrubs, or cutting larger areas of knotweed before planting, may allow woody plants to thrive in, and eventually shade out, knotweed patches.

While both cutting and herbicide injection treatments allowed species richness in the plots to increase, introduced and native plant vigor and moss diversity were higher in the herbicide plots – likely due to the lack of frequent cutting disturbance. Therefore, if primary management goals are to increase plant diversity, herbicide injection may be the better management technique of the two. Hagen and Dunwiddie (2008) found that nontarget vegetation was not affected by glyphosate injection. In addition, herbicide injection may be less expensive than cutting due to high labor costs associated with cutting (J. Grieser, unpublished data). The NYCDEP has a strong preference to avoid chemical treatment, because of the potential for chemical contamination of the water supply. If herbicides are used, they should first be tried on a limited basis with monitoring for herbicide residues and toxic breakdown products. In addition, smaller amounts of glyphosate may be used; Hagen and Dunwiddie (2008) found no difference in knotweed control between 3 ml and 5 ml glyphosate applications. However, if primary management goals are to stabilize stream banks, cutting, possibly combined with woody plantings, may be the better alternative because it alleviates concerns about chemical contamination of the water supply and replaces knotweed with deep-rooted species.

Given that the treatment methods we used require intensive effort and lengthy maintenance to achieve success in reducing knotweed and facilitating colonization by

alternate flora, a strategy other than large scale eradication of knotweed may be appropriate. Such a strategy could include removal of small patches of knotweed (e.g., < 10 m diameter), containment of large patches where expansion would threaten important resources, the creation of openings or corridors of alternate vegetation to promote the colonization of alternate vascular flora and mosses and provide access to the stream for fishing, and the planting of fast-growing, tall, deeply rooted native trees to stabilize stream banks. Effort and maintenance would still be required, but goals could be achieved without massive investment of resources to eradicate large knotweed stands. Remaining knotweed would potentially continue to emit vegetative propagules (stem or rhizome fragments) capable of initiating new stands when dispersed by water, ice, or machinery.

Recommendations for further study include:

- Installing saplings that are substantially taller than the knotweed in the woody plantings plots and then monitoring these plantings to determine whether this method alone can be successful in reducing knotweed stands or at least in stabilizing the stream banks.

- Mowing for an extended period of time in the frequent cutting plots to determine whether knotweed can be eliminated by mowing alone.

- Creating plots with combined treatments, e.g. cutting and woody plantings or cutting and an alternative shading technique.

- Trying a rake-out-the-litter treatment (without anything else) experimentally to see if it encourages plant diversity under the knotweed.

ACKNOWLEDGMENTS

The following people were instrumental in this project: Rich Andreasson, Jenn Grieser, Matt Mariglio, Abbe Martin, and Keith Mellott helped with plot preparation. Knotweed cutting was completed by Sean Antonelli, Amanda Davis, Jenn Grieser, Abbe Martin, and Keith Mellot. Chris Tcimpidis and Jay Truesdell applied the herbicide treatment. Jenn Grieser, Abbe Martin, René VanSchaack, and Laura Weyeneth assisted with study design and data collection. James (Spider) Barbour, Gretchen Stevens, and Mark Folsom (Hudsonia) helped with various aspects of the study. We would also like to thank the Lawrence family for giving us access to their land.

REFERENCES

Baker, R.M. 1988. Mechanical control of Japanese knotweed in an S.S.S.I. Aspects of Applied Biology 16:189-192.

Child, L.W. & M. Wade. 2000. The Japanese Knotweed Manual. Packard Publishing, Chichester, U.K. 123 p.

- Crockett, R.P., P. Burgess, R. Hendrickson, and R. Simpson. 2004. Japanese knotweed (*Polygonum cuspidatum* Sieb. & Zucc.). Monsanto Co., Vancouver, WA and Clark County Weed Management Department, Brush Prairie, WA. Published October 2002, Revised March 1, 2004. 4 p. (March 2004 revision not available online; February 2004 revision available at: <http://www.monsanto.com/ito/pdfs/Knotweed2.03.pdf>)
- Day, L. 2003. Soils report: Greene County Japanese knotweed control project. Report to Greene County Soil and Water Conservation District and New York City Department of Environmental Protection. Delaware County Soil and Water Conservation District. 30 p.
- Hagen, E.N., and P.W. Dunwiddie. 2008. Does stem injection of glyphosate control invasive knotweeds (*Polygonum* spp.)? A comparison of four methods. *Invasive Plant Science and Management* 1: 31-35.
- New York State Department of Environmental Conservation. 2009. Environmental Resource Mapper. Available at: <http://www.dec.ny.gov/imsmaps/ERM/viewer.htm>. Accessed 26-October-2009.
- Schoeneberger, P.J., D.A. Wysocki, E.C. Benham, and W.D. Broderson (eds.). 2002. Field book for describing and sampling soils, Version 2.0. U.S. Natural Resources Conservation Service, National Soil Survey Center, Lincoln, Nebraska.
- Seiger, L.A. 1991. The Nature Conservancy Element Stewardship Abstract for *Polygonum cuspidatum* Sieb. & Zucc. (*Reynoutria japonica* Houtt) (*Fallopia baldschuanica*) [sic]. The Nature Conservancy. <http://tncweeds.ucdavis.edu/esadocs/documnts/polycus.html>
- Seiger, L.A. 1997. The status of *Fallopia japonica* (*Reynoutria japonica*; *Polygonum cuspidatum*) in North America, pages 95-102. In J.H. Brock, M. Wade, P. Pyšek and D. Green (eds.). *Plant Invasions: Studies from North America and Europe*. Backhuys Publishers, Leiden, The Netherlands.
- Seiger, L.A. & H.C. Merchant. 1997. Mechanical control of Japanese knotweed: Effects of cutting regime on rhizomatous reserves. *Natural Areas Journal* 17(4):341-345.
- Talmage, E. & E. Kiviat. 2004. Japanese knotweed and water quality on the Batavia Kill in Greene County, New York: Background information and literature review. Greene County Soil and Water Conservation District and New York City Department of Environmental Protection. 27 p. www.gcsxcd.com/stream/knotweed/reports/litreview/JKandwaterquality.pdf

Table 1. Vascular species associated with *Fallopia japonica* in the four plot types (control, woody planting, herbicide injection, and frequent cutting) in 2003, before treatments. Scientific names follow the New York Flora Atlas (www.newyork.plantatlas.usf.edu); names in brackets = names used by researchers during this study. N = Native, I = Introduced (in New York State). Numbers indicate the number of plots a species was found in for each treatment type (maximum plots = 2 for Control, 3 for Woody Planting [Plant], 2 for Herbicide Injection [Herb], and 2 for Frequent Cutting [Cut] plots).

Scientific Name	Common Name	N/I	2003			
			Control	Plant	Herb	Cut
<i>Alliaria petiolata</i>	Garlic mustard	I	2	3	2	2
<i>Arctium lappa</i>	Greater burdock	I	1			
<i>Calystegia sepium</i>	Hedge bindweed	I				2
<i>Echinocystis lobata</i>	Wild cucumber	N			1	
<i>Erigeron annuus</i>	Daisy fleabane	N				1
<i>Fraxinus americana</i>	White ash	N		1		
<i>Glechoma hederacea</i>	Ground ivy	I	1			1
<i>Hemerocallis</i>	Daylily	I				1
<i>Lonicera X bella</i>	Bell's honeysuckle	I				1
<i>Lysimachia ciliata</i>	Fringed loosestrife	N			1	
<i>Parthenocissus quinquefolia</i>	Virginia creeper	N	1		1	2
<i>Phalaris arundinacea</i>	Reed canary grass	N				1
<i>Prunus</i>	Cherry		1	1	2	1
<i>Saponaria officinalis</i>	Bouncing-bet	I				1
<i>Sambucus nigra</i> L. ssp. <i>canadensis</i> [<i>Sambucus canadensis</i>]	Common elderberry	N				1
<i>Taraxacum officinale</i>	Common dandelion	I	2		1	
<i>Thalictrum</i>	Meadow-rue	N				1
<i>Viola sororia</i>	Common violet	N			1	
<i>Vitis</i>	Grape					1
<i>Zizia aurea</i>	Common Alexanders	N				2
Total number of species			6	3	7	14

Table 2. Vascular species associated with *Fallopia japonica* in the four plot types (control, woody planting, herbicide injection, and frequent cutting) in 2006, post-treatment for woody planting and herbicide injection plots and during treatment for frequent cutting plots. Scientific names follow the New York Flora Atlas (www.newyork.plantatlas.usf.edu); names in brackets = names used by researchers during this study. N = Native, I = Introduced (in New York State). Numbers indicate the number of plots a species was found in for each treatment type (maximum plots = 2 for Control, 3 for Woody Planting [Plant], 2 for Herbicide Injection [Herb], and 2 for Frequent Cutting [Cut] plots).

Scientific Name	Common Name	N/I	2006			
			Control	Plant	Herb	Cut
<i>Acer negundo</i>	Box elder	N				1
<i>Agrostis hyemalis</i>	Winter bentgrass	N			2	
<i>Alliaria petiolata</i>	Garlic mustard	I	2	3	2	1
<i>Allium vineale</i>	Wild garlic	I			1	
<i>Amaranthus</i>	Amaranth				1	
<i>Brassicaceae</i>	Mustard				1	1
<i>Carex</i>	Sedge				2	
<i>Centaurea stoebe</i> L. ssp. <i>micranthos</i> [<i>Centaurea maculosa</i>]	Spotted knapweed	I			1	
<i>Chenopodium album</i>	Lamb's-quarters	I			1	
<i>Cyperus</i>	Flatsedge				1	
<i>Daucus carota</i>	Wild carrot	I			1	1
<i>Digitaria ischaemum</i>	Smooth crabgrass	I			1	
<i>Digitaria sanguinalis</i>	Hairy crabgrass	I				1
<i>Echinochloa crus-galli</i>	Barnyard grass	I			1	
<i>Echinocystis lobata</i>	Wild cucumber	N			1	
<i>Epilobium coloratum</i>	Eastern willow-herb	N			2	
<i>Erechtites hieraciifolius</i>	Fireweed	N			1	
<i>Eupatorium altissimum</i>	Tall boneset	I			1	1
<i>Eupatorium perfoliatum</i>	Common boneset	N			1	
<i>Euthamia graminifolia</i>	Grass-leaved goldenrod	N			1	
<i>Galium</i>	Bedstraw				1	
<i>Galium mollugo</i>	White bedstraw	I				2
<i>Geum aleppicum</i>	Yellow avens	N				1
<i>Gnaphalium uliginosum</i>	Low cudweed	I			1	
<i>Hemerocallis</i>	Daylily	I				1
<i>Hypericum mutilum</i>	Dwarf St. John's-wort	N			1	1
<i>Hypericum perforatum</i>	Common St. John's-wort	I			1	
<i>Impatiens capensis</i>	Spotted jewelweed	N			2	
<i>Juglans cinerea</i>	Butternut	N			1	
<i>Juncus tenuis</i>	Path rush	N			1	
<i>Lapsana communis</i>	Common nipplewort	I			1	
<i>Leersia virginica</i>	Virginia cutgrass	N			1	

Table 2 cont.

Scientific Name	Common Name	N/I	Control	Plant	Herb	Cut
<i>Leonurus marrubiastrum</i>	Lion's tail	I			1	
<i>Linaria</i>	Toadflax	I			1	
<i>Ludwigia palustris</i>	Water purslane	N			1	
<i>Lycopus</i>	Bugleweed				1	
<i>Lysimachia ciliata</i>	Fringed loosestrife	N				1
<i>Lysimachia nummularia</i>	Moneywort	I				1
<i>Mimulus ringens</i>	Square-stemmed monkey-flower	N			1	
<i>Oenothera biennis</i>	Common evening-primrose	N				1
<i>Oxalis stricta</i>	Wood-sorrel	N			2	2
<i>Panicum capillare</i>	Old witch panic-grass	N			1	
<i>Panicum flexile</i>	Wiry panic-grass	N			1	1
<i>Parthenocissus quinquefolia</i>	Virginia creeper	N			1	1
<i>Phalaris arundinacea</i>	Reed canary grass	N			2	1
<i>Plantago lanceolata</i>	English plantain	I			1	1
<i>Plantago major</i>	Common plantain	I			1	1
<i>Poa annua</i>	Annual bluegrass	I				1
Poaceae	Grass				1	1
<i>Polygonum</i> sect. <i>Avicularia</i>	Knotweed					1
<i>Persicaria pensylvanica</i> [<i>Polygonum pensylvanicum</i>]	Pennsylvania smartweed	N			1	
<i>Persicaria punctata</i> [<i>Polygonum punctatum</i>]	Dotted smartweed	N			1	
<i>Persicaria sagittata</i> [<i>Polygonum sagittatum</i>]	Arrowleaf tearthumb	N			1	
<i>Potentilla simplex</i>	Common cinquefoil	N				1
<i>Prunus</i>	Cherry				1	
<i>Prunella vulgaris</i>	Self-heal	I				1
<i>Rhus typhina</i>	Staghorn sumac	N			1	
<i>Rubus allegheniensis</i>	Allegheny blackberry	N			1	
<i>Rumex</i>	Dock				1	
<i>Setaria</i>	Foxtail					1
<i>Solanum dulcamara</i>	Bittersweet nightshade	I			2	
<i>Sonchus asper</i>	Spiny-leaf sowthistle	I			1	
<i>Stellaria</i>	Chickweed	I			1	2
<i>Taraxacum officinale</i>	Common dandelion	I	1		1	2
<i>Trifolium repens</i>	White clover	I				2
Unidentified					2	1

Scientific Name	Common Name	N/I	Control	Plant	Herb	Cut
<i>Verbena hastata</i>	Swamp verbena	N			1	
<i>Verbascum thapsus</i>	Common mullein	I			2	
<i>Veronica</i>	Speedwell	I			1	2
<i>Viola</i>	Violet				2	
<i>Vitis</i>	Grape				1	
Total number of species			2	1	57	30

Table 3. Vascular species associated with *Fallopia japonica* in the four plot types (control, woody planting, herbicide injection, and frequent cutting) in 2009, post-treatment for all plots. Scientific names follow the New York Flora Atlas (www.newyork.plantatlas.usf.edu); names in brackets = names used by researchers during this study.. N = Native, I = Introduced (in New York State). Numbers indicate the number of plots a species was found in for each treatment type (maximum plots = 2 for Control, 3 for Woody Planting [Plant], 2 for Herbicide Injection [Herb], and 2 for Frequent Cutting [Cut] plots).

2009						
Scientific Name	Common Name	N/I	Control	Plant	Herb	Cut
<i>Acer rubrum</i>	Red maple	N			1	
<i>Aegopodium podagraria</i>	Bishops goutweed	I				1
<i>Agrostis hyemalis</i>	Winter bentgrass	N			2	2
<i>Alliaria petiolata</i>	Garlic mustard	I	1		2	2
<i>Arctium lappa</i>	Greater burdock	I			1	1
<i>Berteroa incana</i>	Hoary alyssum	I				1
<i>Bromus inermis</i>	Awnless brome	I				1
<i>Carex</i>	Sedge					2
<i>Clematis virginiana</i>	Virgin's-bower	N				1
<i>Cornus racemosa</i>	Gray dogwood	N			1	
<i>Echinocystis lobata</i>	Wild cucumber	N	1	1	2	
<i>Elymus riparius</i>	River wild-rye	N				1
<i>Epilobium</i>	Willow-herb	N				1
<i>Epilobium coloratum</i>	Eastern willow-herb	N			2	
<i>Erigeron annuus</i>	Daisy fleabane	N			1	
<i>Eupatorium altissimum</i>	Tall boneset	I			1	
<i>Eupatorium purpureum</i>	Joe-pye weed	N			1	
<i>Euthamia graminifolia</i>	Grass-leaved goldenrod	N			2	
<i>Galium mollugo</i>	White bedstraw	I				1
<i>Galium triflorum</i>	Fragrant bedstraw	N			2	
<i>Geum aleppicum</i>	Yellow avens	N				2
<i>Glechoma hederacea</i>	Ground ivy	I				1
<i>Hemerocallis</i>	Daylily	I				1
<i>Hypericum mutilum</i>	Dwarf St. John's-wort	N			2	
<i>Hypericum perforatum</i>	Common St. John's-wort	I				1
<i>Impatiens capensis</i>	Spotted jewelweed	N			2	1
<i>Juncus effusus</i>	Common rush	N			1	1

Table 3 cont.

Scientific Name	Common Name	N/I	Control	Plant	Herb	Cut
<i>Juncus tenuis</i>	Path rush	N			1	1
<i>Lamiaceae</i>	Mint family				1	
<i>Lamium maculatum</i>	White Nancy	I			1	
<i>Lapsana communis</i>	Common nipplewort	I			1	
<i>Leonurus cardiaca</i>	Mother-wort	I				1
<i>Leonurus marrubiastrum</i>	Lion's tail	I			2	
<i>Lepidium</i>	Pepperweed					1
<i>Linaria</i>	Toadflax	I				1
<i>Lysimachia ciliata</i>	Fringed loosestrife	N				2
<i>Lysimachia nummularia</i>	Moneywort	I				1
<i>Mentha arvensis</i>	Wild mint	I				1
<i>Oenothera biennis</i>	Common evening-primrose	N			1	1
<i>Oxalis stricta</i>	Wood-sorrel	N			2	2
<i>Parthenocissus quinquefolia</i>	Virginia creeper	N			2	1
<i>Pastinaca sativa</i>	Wild parsnip	I			2	
<i>Phalaris arundinacea</i>	Reed canary grass	N			1	2
Poaceae	Grass				2	2
<i>Persicaria lapathifolia</i> [<i>Polygonum lapathifolium</i> <i>X persicaria</i>]	Dock-leaved smartweed	I			1	
<i>Persicaria punctata</i> [<i>Polygonum punctatum</i>]	Dotted smartweed	N			1	1
<i>Persicaria sagittata</i> [<i>Polygonum sagittatum</i>]	Arrowleaf tearthumb	N			1	
<i>Potentilla simplex</i>	Common cinquefoil	N				1
<i>Prunus</i>	Cherry				1	1
<i>Prunus serotina</i>	Black cherry	N			1	1
<i>Rosa multiflora</i>	Multiflora rose	I			1	
<i>Rubus occidentalis</i>	Black raspberry	N			2	
<i>Rumex</i>	Dock				1	1
<i>Rumex obtusifolius</i>	Broad-leaf dock	I				1
<i>Sambucus nigra</i> L. ssp. <i>canadensis</i> [<i>Sambucus canadensis</i>]	Common elderberry	N			1	
<i>Scrophularia marilandica</i>	Carpenter's square	N			1	
<i>Secale cereale</i>	Cereal rye	I				1
<i>Solanum dulcamara</i>	Bittersweet nightshade	I			2	
<i>Solidago</i>	Goldenrod	N			1	2
<i>Sonchus arvensis</i>	Field sowthistle	I			1	
<i>Stellaria</i>	Chickweed	I			1	1
<i>Taraxacum officinale</i>	Common dandelion	I	1		2	2

Scientific Name	Common Name	N/I	Control	Plant	Herb	Cut
Unidentified					1	1
<i>Veronica</i>	Speedwell	I			2	1
<i>Verbascum thapsus</i>	Common mullein	I			1	1
<i>Vitis</i>	Grape				1	
Total number of species			3	1	45	42

Table 4. Bryophyte species in *Fallopia japonica* treatment plots, 2006 and 2009. Numbers indicate the number of plots a species was found in for each treatment type (maximum plots = 2).



Scientific Name	2006 Cutting	2009 Herbicide Cutting	
<i>Amblystegium varium</i>			1
<i>Aphanorrhegma serratum</i>	2?		
<i>Atrichum altecristatum</i>		1	2
<i>Barbula unguiculata</i>	1?		1
<i>Brachythecium reflexum</i>		1	
<i>Brachythecium salebrosum</i>		1	1
<i>Bryhnia novae-angliae</i>		2	2
<i>Bryum</i> cf. <i>bicolor</i>		1	
<i>Bryum capillare</i>		1	
<i>Bryum</i> cf. <i>pseudotriquetrum</i>	1	1	
<i>Bryum</i> cf. <i>radiculosum</i>		1	
<i>Bryum</i>		1	
<i>Campylium hispidulum</i>		1	
<i>Campylium radicale</i>		1	
<i>Ceratodon purpureus</i>		2	1
<i>Dicranella heteromalla</i>	1	1	
<i>Ditrichum pusillum</i>		1	
<i>Ditrichum</i>			1
<i>Drepanocladus fluitans</i>		2	1
<i>Ephemerum crassinervium</i>		1	
<i>Eurhynchium pulchellum</i>		1	
<i>Hypnum lindbergii</i>		1	2
<i>Hypnum pallescens</i>	1		
<i>Leptobryum pyriforme</i>		1	
<i>Leptodictyum riparium</i>		1	
<i>Plagiomnium cuspidatum</i>		1	2
<i>Pohlia</i>			1
<i>Polytrichum</i> cf. <i>formosum</i>	1		
<i>Thuidium delicatulum</i>		1	
<i>Weissia controversa</i>		2	
Total number of species	6	23	11


Table 5. Duration of plants in *Fallopia japonica* treatment plots by year.

Plot/Year	Treatment	Annual	Biennial	Perennial Herb	Woody
L1 2003	<i>Control</i>	0	1	2	1
2006		0	1	1	0
2009		1	0	1	0
L6 2003	<i>Control</i>	0	2	1	1
2006		0	1	0	0
2009		0	1	0	0
L2 2003	<i>Woody planting</i>	0	1	1	1
2006		0	1	0	0
2009		1	0	0	0
L4 2003	<i>Woody planting</i>	0	1	0	0
2006		0	1	0	0
2009		0	0	0	0
L5 2003	<i>Woody planting</i>	0	1	0	0
2006		0	1	0	0
2009		0	0	0	0
L3 2003	<i>Herbicide injection</i>	0	1	2	1
2006		4	3	12	5
2009		4	5	14	7
L8 2003	<i>Herbicide injection</i>	1	1	1	2
2006		8	5	20	0
2009		3	4	16	3
L7 2003	<i>Frequent cutting</i>	1	1	4	2
2006		4	2	14	0
2009		2	5	25	0
L9 2003	<i>Frequent cutting</i>	0	1	5	4
2006		0	1	5	1
2009		0	1	13	5

Figure 1. Japanese knotweed study area,
Town of Ashland, Greene County, New York











Figure 8. Box plots of an index of introduced plant vigor (sums of species cover multiplied by maximum height) by treatment type and year




Figure 9. Box plots of average canopy height of *Fallopia japonica* (cm) by treatment type and year




Figure 10. Box plots of percent litter cover (all species) by treatment type and year.




Figure 11. Box plots of average loss of leaf tissue to grazing by treatment type and year




Figure 12. Box plot of average litter depth (all species, cm) in each treatment type in 2009.

