

2.1 Regional Setting

The East Kill watershed is located in the northern Catskill Mountain region of southeast New York State (Fig 2.2.1). The East Kill flows from its headwaters upstream of Lake Capra (upstream of Colgate Lake) to its confluence with the Schoharie Creek in the town of Jewett. The East Kill roughly follows route 23C, paralleling the Schoharie Creek until their convergence. Nearly the entire East Kill watershed is located within the town of Jewett, with small parts entering into Windham, Ashland, and the northern tip of Lexington (Fig 2.1.2). The East Kill begins in East Jewett and ends at Jewett Center. This watershed region is closely surrounded by the steep mountains of the Catskills, specifically the Blackheads.



Figure 2.1.1 East Kill watershed

The entire East Kill watershed lies within the Catskill Park. The Catskill and Adirondack Forest Preserves were established by the NYS Assembly in 1885. An 1894 amendment to the New York State Constitution (now Article 14) directs: "the lands of the State now owned or hereafter acquired, constituting the forest preserve as now fixed by law, shall be forever kept as wild forest lands. They shall not be leased, sold or exchanged, or be taken by any corporation, public or private, nor shall the timber thereon be sold, removed or destroyed (NYS DEC, 2006)."



State Land historical marker

In 1904, the Catskill Park was designated, establishing a boundary or 'blue line' around the Forest Preserve and private land as well. Over the years the Catskill Park grew, and now comprises roughly 700,000 acres, about half of which is public Forest Preserve. The Catskill and Adirondack Parks are nationally unique because they are a checkerboard of public and private land; a grand experiment in how

nature and human society can coexist in a landscape (Catskill Center₁, 2006).

A dominant characteristic of the East Kill watershed's regional setting is its location within the 2,000 square-mile New York City Watershed. The NYC Watershed is the largest unfiltered water supply in the U.S., providing 1.4 billion gallons of clean drinking water each day to over nine million residents in New York City and some smaller municipalities (nearly half the population of New York State) (Catskill Center₂, 2006).



Figure 2.1.2 East Kill watershed towns

The East Kill is a tributary of the Schoharie Creek, which eventually empties into the Schoharie Reservoir where part of the water is transfer through the Shandaken portal to the Ashokan Reservoir. The Ashokan Reservoir supplies approximately 10% of NYC's drinking water. The NYC Department of Environmental Protection (DEP) operates this drinking water supply under a Filtration Avoidance Determination (FAD) issued by the Environmental Protection Agency and the New York State Department of Health. Central to the maintenance of the FAD are a series of partnership programs between NYC and the upstate communities, as well as a set of rules and regulations administered by the DEP. Due to its location within the NYC Watershed, the residents and landowners in the East Kill watershed are subject to the DEP rules and regulations written to protect this watershed.

References

Catskill Center₁, 2006. About the Catskill Region. Available on web: www.catskillcenter.org/region.html (Accessed 12/20/06).

Catskill Center₂, 2006. New York City Watershed. Available on web: http://www.catskillcenter.org/atlas/nycwatershed/nycw_1watershedbasins.htm (Accessed 12/20/06).

NYC DEC, 2006. Catskill Forest Preserve. Available on web: www.dec.state.ny.us/website/dlf/publands/cats/index.html (Accessed 12/20/06).

2.2 Early History of the East Kill Valley

When settlers first arrived in the East Kill Valley, they came upon a rough and wild wilderness. Bears, wolves, and other dangerous wildlife roamed the thickly forested region. It is noted in historical records that many of the first settlers only remained there for a short time before moving away. Therefore, little is known about many of these pioneers. A local account of one of these men from East Jewett further exemplifies some of the difficulties of inhabiting this land. The tale is of Timothy Lockwood who came from Dutchess County to live in the East Kill Valley in 1797. He is said to have retreated almost immediately upon arriving, after being intimidated by the howling wolves (Beach, 1871).

Another man by the name of John Godsell also holds some renown with regards to wolves, but in a different respect. Known as “Uncle John” by the locals, he was an expert trapper. In his time, wolves were a big nuisance to sheep herders who were losing many of their livestock to the carnivores. As a result, New York State encouraged the killing of these wolves by passing laws which allowed towns to offer bounties for them. John Godsell is reported to have killed between 40 and 50 wolves while residing in the East Kill Valley. A story is told of one particular wolf which he tied up and tried to bring to the nearby town of Cairo. His goal was to cross the Cairo line with this captured wolf since the bounty for killing a wolf differed from town to town, and in Cairo it was much higher. However, as a result of Godsell’s cruel treatment, the wolf died before Godsell was able to reach Cairo (Beach, 1871).



The East Jewett Valley, looking east c.1914; from the Tom Hitchcock Collection.

The East Kill is located in the present day town of Jewett. It begins at Colgate Lake in East Jewett and flows west. It roughly follows Rt. 23C, and then turns south to follow Rt. 17. It eventually spills into the Schoharie Creek near Jewett Center where Rt. 17 meets Rt. 23A. The East Kill Valley was

originally part of the town of Windham. It has a history of name changes before it finally became known as Jewett. In 1813 a section of old Windham, encompassing current East Jewett, was broken off. In 1814, the name Hunter was given to this region. The town of Hunter currently borders Jewett, but at the time the two towns were one. It wasn't until 1849 that part of Hunter as well as part of Lexington was sectioned off to form the town of Jewett. The towns name comes from a man named Freeborn G. Jewett. Freeborn was a politician born in Connecticut who held office as U.S. Representative from New York from 1831-1833 and Supreme Court Judge from 1847-1848 (Beach, 1871); (Kestenbaum, 2005).

Industry

In the late 1700's-to-early 1800's, the region thrived from the sale of hemlock bark. Hemlock bark was used for its tannins to tan leather. The tanning industry was becoming quite popular in the Catskill Mountains at that time and the demand for hemlock bark was rising. The local industry later shifted to logging as lumber demands rose. Both these industries proved prosperous for the East Kill Valley. Inevitably the logging and hemlock businesses had negative effects on the environment and sentiment for this subject was not very high as it is today. Eventually, when the land was cleared from all the logging and hemlock harvesting, dairy farms emerged as the dominant economic engine and remained the East Kill Valleys most prosperous industry.

(Beach, 1871).



Slater Farm House (which burned) west of current Farber Farm; from the Alvena Hitchcock Collection.

In 1880 there were 151 farms scattered throughout the area. So many farms in one area can have negative impacts on local streams. For example, many farmers cut their grass fields right up to the streams' edge which creates unstable banks and worsens flood damage. The problem spreads via the stream, which carries these eroded materials creating turbidity and irregular sediment

deposition. Another problem, specifically associated with dairy farms near streams, is the runoff of high levels of fecal matter from cow manure. The presence of fecal coliform bacteria in the water is an indicator of this problem. Fecal coliform bacteria is found in both human and animal waste and makes for unhealthy drinking water. Better understanding and recent technologies have allowed farmers to employ more environmentally friendly ways of managing their farms than practiced in the past.

Due to economic pressures many of the local farms have gone out of business. J.J. Farber Farm is one of the only remaining farms in the East Kill Valley. Located at the site of former Slater Farm, they have approximately fifty beef cows. The East Kill Bridge (Farber Farm Bridge) stretches across the East Kill Creek here, connecting the farmhouse to more of the farms' cattle fields. This bridge was built in 1932 and used to be known as Slater Bridge, when Slater Farm was running (Historic Bridges of the United States, 2006). GCSWCD, in collaboration with NYCDEP, has been involved in an ongoing project to stabilize the creek at this location and alleviate localized flooding.



Bridge at Slater Farm, current Farber Farm; from the Alvena Hitchcock Collection.

Today

The East Kill Valley never experienced quite as big of a boom in tourism as has been characteristic of surrounding areas in the Catskills, such as Windham and Hunter. It has remained in many ways as it began, a peaceful area offering a simple lifestyle, known for its wild forests and the East Kill Creek. It is currently one of the least populated towns in Greene County. The undeveloped nature of the area is largely due to the fact that New York State owns much of the forests and mountain peaks. The state maintains these areas as protected wilderness, preserving the natural landscape and maintaining hiking trails. Many of its residents still thrive off agriculture and agroforestry. The number of farms has decreased dramatically since its early days, with current farming practices and techniques becoming much more environmentally stable. The area offers a vacation spot

for tourists who are looking for serenity, peace, and quiet in a less commercialized location of the Catskills (Greene County Planning and Economic Development, 2004). Three popular hiking destinations, Blackhead, Blackdome, and Thomas Cole Mountain, are all located in the northeastern part of the valley.



From left: Blackhead, Blackdome, and Thomas Cole mountains; from www.catskillguide.com

References

- Beach, H. N. 1871. Early Settlers of East Jewett. Catskill Examiner. Available on Web: http://www.rootsweb.com/~nygreen2/early_settlers_of_east_jewett.htm (Accessed 11/29/06).
- Greene County Planning and Economic Development. 2004. Community Profile, Town of Jewett. Available on Web: <http://www.greeneplanning.com/greene/communityprofiles.htm> (Accessed 11/29/06).
- Historic Bridges of the United States. 2006. East Kill Bridge (#1045150). James Baughn and Contributors. Available on Web: <http://bridges.midwestplaces.com/ny/greene/1045150/> (Accessed 11/29/06).
- Kestenbaum, Lawrence. 2005. Index to Politicians: Jewett. The Political Graveyard. Available on Web: <http://politicalgraveyard.com/bio/jewett.html> (Accessed 11/29/06).

2.3 Physical Geography of the East Kill Watershed

Physical geography encompasses the physical elements and processes that comprise the earth's surface features and associated processes. These processes include: energy, air, water, weather, climate, landforms, soils, animals, plants, and the Earth itself. The study of physical geography attempts to explain the geographic patterns of climate, vegetation, soils, hydrology, and landforms, and the physical environments that result from their interactions.




Blackheads-East Kill/Batavia Kill valley divide in background.

The East watershed is located in the Appalachian Plateau physiographic province (Figure 2.3.1). The erosional characteristics of the sedimentary rock formations of the Appalachian Mountains are responsible for the characteristic valley and ridge topography of the Catskills. Durable layers of sandstone and conglomerate form ridges and less resistant limestone and shale underlie the East Kill valley as it winds its way to the Schoharie Creek. During the height of glaciations, the East Kill watershed was covered by an ice sheet up to a mile thick. Upon retreat, the ice sheet left a layer of unsorted and unconsolidated glacial debris, glacial till, ranging from clay particles to huge boulders. Following the retreat of the ice sheet, the landscape was covered with glacial tills and bedrock and was wiped clean of plants and animals, leaving a clean slate for the migration and colonization of the modern plant and animal communities. Today, the East Kill watershed lies within the Northeastern Highlands ecoregion. This ecoregion is characterized by nutrient poor soils blanketed by northern hardwood and spruce fir forests.



Figure 2.3.1. Physiographic Regions of the United States, including the Appalachian Plateau (NASA Earth Observing System (EOS) Goddard Program Office).

Elevations in the watershed vary from a high of approximately 3,948 feet above sea level on the top of Black Dome Mountain in Southeast Jewett, to a low point of 1,400 feet above sea level at the Schoharie Creek/East Kill confluence. The average elevation of the watershed is approximately 2,674 feet above sea level. Studies indicated that the temperature drops approximately 3.0° F per 1000' of elevation (Thaler, 1996). The East Kill starts as a mountainous stream dropping approximately 780 feet in its first mile (Eastern Jewett), then reducing in slope to approximately 200 feet in the next mile, and then to an average slope of 42 feet per mile until it reaches its confluence with the Schoharie Creek. The more notable high peaks (>3,500') that form the East Kill watershed boundary with the Batavia Kill Thomas Cole (3,940'), Black Dome (3,980') and Blackhead (3,940') (Figure 2.3.2).



The East Kill flows west through the Town of Jewett before entering the Schoharie Creek near the intersection of County Route 17 and State Route 23a. The watershed contains approximately 94 miles of stream, including Halsey Brook near Colgate Lake and Roaring Brook, just upstream of where EK crosses Mill Hollow Rd.

Climate

The climate of the East Kill basin is primarily driven by the humid continental type, which dominates the northeastern United States. The average annual temperature for the area is 44.8° F and the area typically receives approximately 41” of rain/year (Table 2.3.1 and Figure 2.3.2). Due to up-sloping and down-sloping, the character of the mountaintop topography can affect the climate of the basin. Up-sloping occurs when air is lifted up over the mountains, the air expands, cooling and condensing into moisture, which takes the form of clouds and precipitation (Thaler, 1996). Down-sloping occurs when air sinking within a dome of high pressure or air that is forced downslope of a mountain range, warms up and loses moisture, as is shown by a drop in relative humidity (Thaler, 1996). These weather phenomena can be responsible for differences in cloud cover and precipitation between the Catskills and the surrounding area, and helps to explain the sometimes drastic variations in rainfall between Catskill basins (Figure 2.3.3).

Table 2.3.1. Average annual temperature, precipitation, snow fall and winter and summer temperatures for Windham, NY from the period 1961-1990 (Thaler, 1996).	
Average Annual Precipitation	41”
Average Annual Temperature	44.8 ° F
Average Winter Temperature	25.6 ° F
Average Summer Temperature	64.7 ° F
Seasonal Snowfall	60”





Figure 2.3.3. Radar showing the rainfall intensity that caused the flooding in the Western Catskills in June 2006. The isolated pockets of heavy rain (dark red) within individual Valleys help explain why flood damages can be so dramatic from one basin to the next (National Weather Service Forecasting Office).



Global Climate Change Effects on the Watershed

Global warming will impact the Schoharie basin in coming years. Greenhouse gases are trapping energy in our atmosphere that would normally be lost to space and causing global temperatures to rise. This warming is a natural phenomenon that provides enough heat to allow humans to thrive on earth, but the burning of fossil fuels, and the atmospheric concentration of other gases such as methane, has dramatically increased the rate of warming (Figure 2.3.4). Based on local data collected between 1952 and 2005, researchers have concluded that a broad general pattern of warming air temperatures, increased precipitation, increased stream runoff and increased potential evapotranspiration has occurred in the Catskills region (Burns et al., 2007). In coming years, there is no doubt that the effects of global warming will impact management decisions in the East Kill watershed.




Figure 2.3.4. Millennial northern hemisphere temperature reconstruction, based upon ice core data, relative to actual temperatures recorded from 1902 through 1999. Despite large variation, the recent trend of rapid heating in the industrial era is apparent (National Climatic Data Center adapted from Mann et al., 1999).

Temperature increases will have effects on food production, plants, wildlife, invasive species, flooding, drought, snowfall and the economy. Based upon current climatic trends,

our climate may migrate to the extent that by the end of the century, summers in upstate New York may feel like Virginia (Figure 2.3.5) (Frumhoff et al., 2006). This climatic migration will have deleterious effects on plant and animal life, allowing new warmer climate species to thrive at the expense of our traditional plants and animals. The number of snow-covered days across the Northeast has already decreased, as less precipitation falls as snow and more as rain, and as warmer temperatures melt the snow more quickly. By the end of the century, the southern and western parts of the Northeast could experience as few as 5 to 10 snow-covered days in winter, compared with 10 to 45 days historically (Frumhoff et al., 2006). Decreased snowfall and increased rainfall would have negative effects on stream flows and the economy of the Catskills.

With the lack of snow fall, streams and groundwater will not receive a slow sustaining release of water through the winter and spring. Replacing the slow release will be more intense storms, which will sporadically dump large quantities of water into the system potentially causing damaging flooding (Figure 2.3.6). However, streams will return to base flow relatively quickly once the rain stops. Modeling predictions indicate that in the next century we will see more extreme stream flows that will cause streams to flow higher in winter, likely increasing flood risk, and lower in summer, exacerbating drought (Frumhoff et al., 2006). Changing the dynamic of the hydrologic cycle would also impact the NYC water supply system, forcing potential changes in operational measures.



Figure 2.3.5. Projected climate “migrations” for Upstate, NY based on average summer heat index, under the lower (yellow)- and higher-emissions (rust) scenarios. Based on the average of the GFDL, HadCM3 and PCM model projections (Frumhoff et al., 2006).




Figure 2.3.6. Projected increases in three indices of extreme precipitation: (1) precipitation intensity, (2) number of days per year with more than two inches of rain, and (3) maximum amount of precipitation to fall during a five day period each year (Frumhoff et al., 2006).

Because we don't have a clear understanding of all of the coming impacts of climate change, stream managers need to employ the "no-regrets policy" with regard to their current management actions and policies. The no-regrets policy is the recognition that lack of certainty regarding a threat or risk should not be used as an excuse for not taking action to avert that threat, that delaying action until there is compelling evidence of harm will often mean that it is then too costly or impossible to avert the threat. Stream managers—including streamside landowners-- will need a basic understanding of how streams are formed and evolve to effectively adapt to coming changes. They will need to anticipate and compare the consequences of different management options, and will need to act conservatively: oversizing culverts and bridge spans, leaving larger buffers of undisturbed streamside vegetation, and consider limiting new development of infrastructure or personal property in areas where conditions indicate a high risk of the stream channel shifting across the floodplain. The humid continental climate has been an unquestionable asset to the historical development of the East Kill basin and its many occupants and uses. With proper planning and implementation of the no-regrets policy, undoubtedly, the climate will continue its important role in East Kill basin life.

References

- Burns, D.A., Klaus, J. and McHale, M.R. 2007. Recent Climate trends and implications for water resources in the Catskill Mountain region, New York, USA. *Journal of Hydrology* (2007), [doi:10.1016/j.jhydrol.2006.12.019](https://doi.org/10.1016/j.jhydrol.2006.12.019).

Frumhoff, P., McCarthy, J., Melillo, J., Moser, S., Wuebbles, D. 2006. Climate Change in the U.S. Northeast: A Report of the Northeast Climate Impacts Assessment. Union of Concerned Scientists: Cambridge, MA. Available on web: <http://www.northeastclimateimpacts.org>.

Thaler, J.S. 1996. Catskill Weather. Purple Mountain Press, Fleischmanns, NY.

2.4 Hydrology and Flood History

Introduction

Hydrology is the study of the properties, distribution, and effects of water on the earth's surface, in the soil and underlying rocks (groundwater), and in the atmosphere. The *hydrologic cycle* includes all of the ways in which water cycles from the landscape (both underground and in streams and water bodies) to the atmosphere (as water vapor and clouds) and back (as snow, rain and other forms of precipitation) (Figure 2.4.1). Understanding the hydrology of the East Kill will assist us with making land use decisions in the basin that work within the constraints of the hydrologic cycle and won't exacerbate flooding or cause water quality impairment.




Figure 2.4.1. The Hydrologic (water) Cycle
(<http://ga.water.usgs.gov/edu/watercyclesummary.html>).

Water flowing through the East Kill reflects the integrated effects of all watershed characteristics that influence the hydrologic cycle. Characteristics include climate of the drainage basin (type and distribution patterns of precipitation and temperature regime), geology and land use/cover (permeable or impermeable surfaces and materials affecting timing and amount of infiltration and runoff, and human-built drainage systems), and vegetation (uptake of water by plants, protection against erosion, and influence on infiltration

rates). These factors affect timing and amount of stream flow, referred to as the stream's *hydrologic regime*. For example, a stream with an urbanized watershed where water will run off the hardened surfaces directly into the stream will have higher peak discharges following storms than a watershed, such as the East Kill, which is predominantly forested and will infiltrate a higher percentage of rain water before it reaches the stream and release it more slowly over time. Understanding the hydrology of a drainage basin is important to the stream manager because stream flow patterns affect aquatic habitat, flood behavior, recreational use, and water supply and quality.

East Kill Basics

Encompassing approximately 37 square miles of watershed area almost exclusively in the Town of Jewett, the East Kill watershed is typical of main headwaters tributaries to the Schoharie Creek in that it is a long, narrow watershed running east to west. This drainage pattern is controlled by the steep topography, formed in large part during the last period of glacial activity. Streams in the East Kill valley are primarily perennial streams, that is, they flow year-round except in smaller headwater streams or in extreme drought conditions.

The East Kill watershed averages approximately 44 inches of precipitation per year that often comes in dramatic summer downbursts, remnants of autumn hurricanes, or late winter rain-on-snow events. Average slope of the watershed is 17.3% (watershed elevation drops 17.3' feet for every 100 feet horizontal distance). *Drainage density*, or how much stream length is available to carry water off the landscape is slightly higher than average for the Catskills, at 0.0016m/m . Given the average drainage density, combined with steep mountainous slopes, and high precipitation, the East Kill stream system is relatively *flashy*, that is, stream levels rise and fall quickly in response to storm events. This flashiness is somewhat mitigated by heavy forest cover throughout much of the watershed. Therefore, efforts to protect upland, as well as riparian, forest are important to reduce flooding.

Stream flow Primer

There are two general categories of streamflow: storm flow (also called flood flow) and base flow, between which streams fluctuate over time. Storm flow fills the stream channel in direct response to precipitation (rain or snow) or snowmelt, whereas base flow is

primarily groundwater fed and sustains streamflow between storms and during subfreezing or drought periods. A large portion of storm flow is made up of *overland flow*, runoff that occurs over and just below the soil surface during a rain or snowmelt event. This surface runoff appears in the stream relatively quickly and recedes soon after the event. The role of overland flow in East Kill watershed is variable, depending upon time of year and severity of storms or snowmelt events. In general, higher streamflows are more common during spring due to rain, snowmelt and combination events, and during hurricane season in the fall. During summer months, actively growing vegetation on the landscape draws vast amounts of water from the soil through *evapotranspiration*. This demand for groundwater by vegetation can significantly delay and reduce the amount of runoff reaching streams during a rain storm. During winter months, precipitation is held in the landscape as snow and ice, so precipitation events do not generally result in significant runoff to streams. However, frozen ground may increase the amount of overland flow resulting from a rain storm if the air temperature is above freezing, particularly in spring on north facing slopes.

Subsurface storm flow, or *interflow*, comes from rain or snow melt that infiltrates the soil and runs down slope through the ground. Infiltrated water can flow rapidly through highly permeable portions of the soil or displace existing water into a channel by “pushing” it from behind. In the East Kill valley, subsurface flow can occur fairly rapidly along layers of essentially impermeable glacial lake silt/clay deposits. Subsurface storm flow shows up in the stream following overland flow, as stream flow declines back to base flow conditions.

Base flow consists of water that infiltrates into the ground during and after a rain storm, sustaining streamflow during dry periods and between storm flows. The source of base flow is groundwater that flows through unsaturated and saturated soils and cracks or layers in bedrock adjacent to the stream. In this way streams can sustain flow for weeks or months between precipitation events and through the winter when the ground surface and all precipitation is otherwise frozen. Stable-temperature groundwater inputs is what enables fish and other aquatic life to survive in streams year-round.

Hydrologists use a hydrograph of a stream to illustrate specific hydrologic measurements, such as water level, discharge or velocity, over a period of time. A *stream gage* is necessary to monitor stream discharge and develop a hydrograph. The United States

Geological Survey (USGS) maintains a continuously recording stream gage on the East Kill near Jewett Center (established 1996, drainage area 35.6 mi², USGS ID# 01349700). Prior to 1996, a crest stage gage was maintained starting in 1929. All gage information is available online at the USGS website: http://waterdata.usgs.gov/ny/nwis/uv/?site_no=01349700. This gage measures the *stage*, or height, of the water surface at a specific location, typically updating the measurement every 15 minutes. By knowing the stage, we can calculate the magnitude of the *discharge*, or volume of water flowing by that point using a relationship developed by USGS called a *rating curve*. Using this rating curve, the magnitude of flow in East Kill at the gage location can be determined at any time just by knowing current stage, or flow can be predicted for any other stage of interest. Additionally, we can use the historic record of constantly changing stage values to construct a picture of stream response to rain storms, snow melt or extended periods of drought, to analyze seasonal patterns or flood characteristics.

The East Kill gage has a long enough period of record to prepare a hydrograph for the stream (Figure 2.4.2). Each spike on the graph represents a peak in stream flow (and stage) in response to rain storms. Stream level rises (called the “rising limb” of the hydrograph) and falls as the flood recedes (called the “falling (or receding) limb” of the hydrograph). We can analyze long time periods to see seasonal trends or long-term averages for the entire length (period) of gage record. We can see the record for the gage shows higher flows in fall (hurricane season) compared to winter (water held in ice and snow), and higher flows in spring (snow and ice melt, with rain-on-snow events) compared to summer (drought conditions with vegetation using a lot of water). The highest flows of the year are generally associated with the hurricane season in the fall, followed by winter and spring snowmelt or rain-on-snow events. Overland flow accounts for most of water that causes the sharp peaks in the hydrograph.

Streamflow always rises and peaks following the height of a precipitation event because it takes time for water to hit the ground and run off to the stream (this is known as lag time). Knowing storm timing, we could also calculate lag time for East Kill at the gage location for particular storms or types of storms, and determine how the stream responds to storms both in timing and flood recession.




Figure 2.4.2. This hydrograph represents the daily average flow from 12/96 through 12/06.

The hydrograph of April, 2005 illustrates the effects of a spring storm on top of snow (Figure 2.4.3). The East Kill rose quickly from the precipitation from a daily average of 411 CFS to 2,290 CFS in 24-hours. The recession took longer than a large summer storm due to the vegetation still being dormant, or just emerging, and the snow pack.




Figure 2.4.3. This hydrograph represents the daily average flow for April, 2005, including a large rain on snow precipitation event.

East Kill Flood History

As a mountain stream the East Kill rises quickly as precipitation falls. Between 1951 and 2006, the East Kill at Jewett Center has gone over 10,000 CFS five times (Figure 2.4.4). For the period of record, the annual average peak flow was 5,816 CFS. The peak flow in any given year is not necessarily a significant/damaging storm event, but could be a dramatic increase in flow following drought. To put this in perspective, the flood of record in 1996 pushed the East Kill to its highest stage at 17' (max flow of 13,500 CFS daily average) reaching its 100 year flood stage (as it did the Schoharie). After this flood \$15.2 million of federal and state funding was distributed amongst 377 municipalities to help repair damage. Of this, the town of Jewett received \$52,135. FEMA estimated that approximately \$102 million worth of damage had occurred state wide during the flood (New York State, 1996). The flooding in 2000 was much less than 1996 (stage height 10.5' / 4,140 CFS), but still brought comparable damage, with \$12.7 million being released to 206 municipalities across NY. Greene County received \$176,596.23. Only three towns within Greene County received additional funding following this flood event. The three received \$44,320.79 in total and Jewett was one of them, receiving \$11,041.23 of the total (New York State, 2000).




Figure 2.4.4. Annual peak flows for the period 1951 through 2006.

Flood events have not been as well documented for the East Kill Valley as other areas due to its relatively small population. However, considering its location and relationship with the Schoharie Creek, it is safe to assume that flooding has historically been a problem in the East Kill, as it has been throughout the Schoharie Valley. North facing slopes of the East Kill valley receive little sun exposure compared to south facing slopes. As a result, half of the valley retains a snow pack well into the spring when rain on snow events can cause dramatic spring flooding. Tropical storms and hurricanes in the late summer and early fall also trigger flooding in the valley. Through analysis of the long-term flood records provided by the USGS, the town, its residents and resource managers can begin to better understand the cause/effect of various precipitation amounts on flooding.

Flood frequency distributions show flood magnitude for various degrees of probability (or percent likelihood). This value is most often converted to a number of years, called the “recurrence interval” (RI) or “return period”. For example, the flood with 20% chance of occurring or being exceeded in any single year corresponds to what is commonly referred to as a “5-year flood” (each of these values is the inverse of the other - just divide 1 by % probability to get RI in years, or divide 1 by RI in years to get % probability). This simply means that on average, for the period of record (the very long term), this magnitude flood will occur about once every 5 years. This probability is purely statistical; probability remains the same year to year over time for a particular size flood to occur, though the actual distribution of flood events in time is not regular; many years may go by without a certain magnitude flood, or it may occur several times in a single year. As another interesting characteristic of flood frequency distributions, the 5-year flood may not occur the “right” number of times in a certain period of record. For example, we might expect to see about 2 “5-year floods” for every 10 years of record, but any particular 10 year period may contain greater or fewer of this size flood.

Because the flood frequency curve is not linear, that is, the shape of the curve doesn’t progress along a steady line, we can’t simply divide up the floods in a record in rank them in order. For example, in a 10 year record, the largest flood is not necessarily a 10-year flood, even though that flood only occurred once in that ten year record. The length of gaging records is typically short compared to long-term history, on the order of 10-30 years, whereas 200-300 years might give a better picture of how often the range of floods may occur.

Therefore, we need to fit some other probability to the floods we do see, based on their magnitude in relation to the other floods in the record, and the average shape of distributions for very long-term records – so individual floods can be plotted where they belong in a more accurate risk of occurrence. Floods recorded at the Esopus, Bushnellsville and Prattsville gages that exceed a 5-year recurrence interval provide an example of distribution of medium to large floods over time (Table 2.4.1).

Table 2.4.1. Flood Flows at Nearby Gages that Exceed Five Year Recurrence Intervals (Flood frequency statistics based on recorded peak flows through 1997. Esopus Creek at Allaben, NY: 5 yr RI flood:~6,500 cfs 10 yr RI flood: ~9,500 cfs Bushnellsville Creek at Shandaken, NY 5 yr RI flood:~800 cfs 10 yr RI flood: ~1,000 cfs Schoharie Creek at Prattsville, NY: 5 yr RI flood: ~24,000 cfs 10 yr RI flood: ~33,000 cfs.).	
Esopus Creek at Allaben, NY	
Date	Flood Discharge (cfs)
3/30/51	20,000
7/28/69	7,870
3/21/80	15,900
2/20/81	6,540
4/5/84	8,470
4/4/87	16,100
1/19/96	15,000
9/18/04	6,700
4/02/05	20,400
Bushnellsville Creek at Shandaken, NY	
Date	Flood Discharge (cfs)
11/25/50	1,350
10/15/55	1,830
3/21/80	845
4/5/84	896
4/4/87	1,000
1/19/96	996
9/18/04	No data available
4/02/05	No data available
Schoharie Creek at Prattsville, NY	
Date	Flood Discharge (cfs)
Sep. 30, 1924	29,000
Nov. 16, 1926	42,300
Aug. 24, 1933	39,000
Mar. 03, 1934	50,002
Jul. 08, 1935	27,400
Mar. 18, 1936	38,500
Feb. 22, 1937	29,800
Sep. 21, 1938	45,000
Nov. 25, 1950	49,500

<i>Dec. 11, 1952</i>	28,200
<i>Aug. 13, 1955</i>	25,100
<i>Oct. 16, 1955</i>	51,600
<i>Dec. 21, 1957</i>	31,000
<i>Sep. 12, 1960</i>	49,900
<i>Jun. 22, 1972</i>	27,400
<i>Dec. 21, 1973</i>	24,900
<i>Dec. 08, 1974</i>	24,800
<i>Jan. 09, 1978</i>	30,600
<i>Mar. 21, 1980</i>	39,600
<i>Apr. 05, 1984</i>	29,500
<i>Apr. 04, 1987</i>	47,600
<i>Jan. 19, 1996</i>	52,800
<i>Sep. 16, 1999</i>	42,800
<i>Sep. 18, 2004</i>	26,500
<i>Apr. 2, 2005</i>	42,500

However, recurrence interval can be misleading if a flood of a certain size is expected to occur at regular intervals. For example, during the 1980s four floods exceeding the “5-year event” occurred within a seven-year span on the Esopus, while there were no such events during the entire decade of the 1970s. On the Schoharie Creek in the 1930s, there were significant floods six years in a row, with two greater than the 25-year event – the size flood for which most NYS and county bridges are designed. By contrast, there were no such events during the entire decade of the 1940s.

Flooding occurs in response to excessive runoff associated with spring snowmelt, summer thunderstorms, fall hurricanes, and winter rain-on-snow events. Five of the seven major floods recorded at the Esopus Creek at Allaben station occurred in late winter/early spring and are presumably associated with major snowmelt events from either spring thaw or rain-on-snow events. The largest recorded flood is a spring runoff event. A summer flood in 1969 and the flood of January 1996 are the two other large floods recorded at the gage. Three of the six major floods recorded at the Bushnellsville gage occurred during the spring and are coincident with three of the Esopus events, showing some comparison can be made between nearby streams. Conversely, weather in the Catskills can produce localized historically significant flood events such that a peak event may not be recorded at each gage for the same time period or storm event. Significantly, we can see that 10 of 25 events at Schoharie Creek occurred during hurricane season (late summer to late fall), 13 occurred

during winter and spring, and only 2 occurred during summer. The January 1996 flood was approximately a 10-year recurrence interval flood on the Bushnellsville Creek, less than a 40 year event at Esopus Creek, and the “flood of record” at the Schoharie Creek. This shows that between-stream comparisons are not always perfect. This is especially so with summer thunderstorms, where highly localized storm cells can produce 10 or more inches of rain in one watershed, and only a few inches in an adjacent watershed for the same storm. Summer peaks shown in Table 2.4.1 do not overlap between any of the three sites.

From review of available data we can generalize that most bankfull and greater events will occur in late winter/spring as the result of thaws and major rain-on-snow events. This is in large part due to landscape storage of available water as snow and ice, reduced infiltration capacity if the ground is still frozen (or partially so), and minimal evapotranspiration from vegetation, which would otherwise route moisture back into the atmosphere. Other major floods can be expected during hurricane and tropical storm season in the late summer and fall, particularly as vegetation enters the dormant season and demand for water in the landscape drops off.

The 1990s were generally a time of moderate flood events in the vicinity of the East Kill, with the exception of the winter flood of January 19, 1996, which was similar in scale to April 1987. Tropical Storm Floyd flood (September 1999) was typical of tropical storm events and sometimes uneven distribution of precipitation associated with those storms. While flooding in Esopus drainages was typically less than a 5-year event, several drainages in bordering Schoharie system had over a foot of precipitation in 24 hours with flooding that exceeded the 10-year event discharge.

The years 2000 – 2002 were characterized by droughty conditions with intervening wet conditions. High water events were typically limited to bankfull (or smaller) events. 2003 was an unusually wet year, with several larger than bankfull events occurring during the summer. Predicting precisely when the next 5-year (or greater) flood will occur in the East Kill is impossible – the probability for a large flood, or a flood of any particular size, is the same each year – though weather and storm patterns can be used to anticipate conditions for a few months out, and seasonal patterns are generally reliable. The last really large flood

was in April, 2005, but the probability is high that, when the next flood occurs, late winter/early spring during snowmelt/rainy season will be prime time.

Implications of East Kill Flooding

The unique hydrology of the East Kill has consequences for how the stream corridor should be managed. Flood history and dynamics play a large role in determining the shape, or morphology, of stream channels and the hazards associated with land uses on the banks and in the floodplain. For example, applications for stream disturbance permits (from NYS DEC) typically increase following floods, as landowners and municipalities attempt to repair damage caused by floods. If we want to minimize their impact on property, infrastructure and other damages or inconvenience, it is critical that we understand and plan for flooding behavior. Historically, this “planning” has emphasized attempts to constrain and control stream channels, rather than working with processes we can measure and, to some extent, predict. The results are often costly, and sometimes catastrophic, such as when berms or levees fail, or bridges wash out. These “control” approaches typically result in ongoing maintenance costs that can draw valuable community resources away from other projects. With a better understanding of stream and floodplain processes, we can reduce these costs. For more information, see Section 3.2, Introduction to Stream Processes.

2.5 Upper Schoharie Watershed Geology (including East Kill)

(Note: this is an adaptation of the Upper Esopus Creek Management Plan geology section)

Introduction

Water flows across the landscape and sculpts the watershed. The geology (the earth material) of the watershed helps determine the nature of the streams, influences the stream's water quality and the way the landscape erodes (Photo 2.5.1). In the Catskill Mountains, geology is the primary influence on water quality. Jill Schneiderman, a professor of geology at Vassar College, notes in her book, *The Earth*



Photo 2.5.1. Streambank erosion into glacial lodgement till along East Kill, tributary to Schoharie Creek.

Around Us: Maintaining a Livable Planet, that the bedrock and glacial sediments of the Catskills provide excellent filtration for maintaining high water quality (Schneiderman, 2003). However the geology also periodically degrades the water quality. Where the stream erodes into very fine-grained (silt and clay) glacial deposits the water will become brown with the suspended sediment. This Section presents basic background information on Catskill and Upper Schoharie watershed geology, and discusses some of the important implications of geology on stream management. The intent is to provide just enough information to describe the geologic setting and history of the Upper Schoharie watershed. Specific recommendations pertaining to further characterization are presented at the end of this Section. References are provided for the reader interested in obtaining more detail on the geology of this region.

Streams and glaciers sculpted these mountains out of a plateau of rock that formed from ancient rivers. That is essentially the geologic story of the Northeastern Catskill Mountains. These mountains and their river valleys are the ongoing result of water interacting with landscape geology under the force of gravity over millions of years.

Knowing the geology of the landscape and stream corridor helps stream managers understand important conditions that control the stream's work (moving water and sediment out of the watershed), as well as significantly influencing water quality.

The nature of the bedrock – its composition and structure – determines how the stream valleys form and what the sediment will be like. Upper Schoharie Creek and its tributaries drain much of the northeast Catskill Mountains. These mountains are composed of sedimentary rock. The broken bits of this rock, formed from layers of ancient river sediment, is the source of almost all of the stream sediment you see today - from clay to boulders. The layered reddish clays exposed in stream banks are



Photo 2.5.2. Streambank/bed with alluvial and non-alluvial sediment

the legacy of ancient lake sediments eroded from the red siltstones and shales that often form the mountain slopes and the cobbles and boulders eroded from the thick-bedded sandstones that form the mountain cliffs (Photo 2.5.2). Much of this sediment that the stream is currently conveying was deposited during the most recent ice ages of 12,000 – 25,000 years ago, when the Catskills were mostly occupied by ice or the melt-water streams and lakes that followed the ice's retreat. The Schoharie Creek and all the streams that feed it water and sediment have inherited this geologic framework.

The geology of the Upper Schoharie Creek valley is typical of the complex geologic conditions that prevail in the tributaries as previously documented in the Batavia Kill (GCSWCD, 2003) and West Kill (GCSWCD, 2005) Stream Management Plans and in the adjacent Esopus Creek basin to the south, as documented in the Upper Esopus Creek Stream Management Plan (CCE, 2007). The bedrock geology is straightforward, while the glacial geology provides the complexity that makes these basins unique in the Catskills.

Bedrock Geology

The bedrock geology of the Catskill Mountains and Upper Schoharie watershed exerts considerable control on the character of its valley slopes and streams (Figure 2.5.1; Photo 2.5.3). The gently sloping sedimentary rock, primarily composed of alternating layers of sandstone and siltstone/shales, creates the characteristic Catskill stepped topography. The sandstones form the cliffs while the more easily erodible siltstones/shales tend to form the slopes. The mountain tops tend to be formed of conglomerate (a gravelly sandstone). The sediments that form the middle-to-late Devonian (390 to 360 million years ago) bedrock are interpreted to be deposits of a vast deltaic river system, often called the “Catskill Delta” deposits that drained the ancient high peaks of the Taconic mountain range (Isachsen et al., 2000). Titus (1998) compared it to the Bangladesh river complex draining the Himalayas. The sandstone and conglomerate are made up of river channel sand and gravel, while the siltstones and shales are overbank and shallow fresh water silts and clays.



Photo 2.5.3. Bedrock exposed along Schoharie Creek

The Catskill Delta deposits were buried beneath younger sediments, and then uplifted as a plateau. Prior to and during the uplift, intersecting sets of vertical fractures formed in the Catskill rock. The following eras eroded away the overlying rock, and streams incised multiple channels into the slowly rising plateau. The following two publications are recommended for further detail on the Catskill bedrock geology: *Geology of New York: A simplified account* (Isachsen, et al., 2000) and *The Catskills: A Geological Guide* (Titus, 1998).

Fisher, et al. (1970) mapped the bedrock of the area as part of the New York State Geological Survey Map and Chart Series. The mapped geologic formations that make up most of the watershed are the Oneonta and Walton formations comprising sandstones, shales,

and mudstones (Photo 2.5.3). Around the Schoharie Reservoir a similar sequence of rocks comprise the Genesee Group and the Moscow formation. The Moscow formation is the rock that hosts the famous Gilboa forest fossils (VanAller Hernick, 1996).




Figure 2.5.1. Bedrock geology of the Schoharie Basin.

The orientation of the stream valley is important, influencing the microclimate, average depth of snowpack and local hydrological regime in many ways. The Schoharie Creek watershed is uniquely oriented for Catskill drainage basins, with drainage to the west and north rather than to the southeast or southwest typical of the other principal watersheds.

Bedrock fracture orientations in the area are generally consistent with the overall trend of two sets for the central to northeastern Catskill Mountains. The dominant high-angle (near vertical) fracture sets are oriented NE and NW, influencing drainage pattern development.

Modern stream deposits in the Catskill Mountains are principally derived from erosion of the well-bedded sedimentary Catskill bedrock. As a result, stream clasts (sediment particles and classes) have a low sphericity (“roundness”), typically forming platy or disk-like particle shapes. This platy shape affects the stability of the streambed in a number of ways. First, it allows the particles to *imbricate*, or stack up at an angle,



Photo 2.5.4. Example of imbricated Catskill stream sediment

forming an overlapping pattern like fish scales or roof shingles (Photo 2.5.4). Imbricated streambeds are thus generally more stable or “locked up”, and all other things being equal, generally require a larger flow to mobilize the bed material than nonimbricated beds. However this same platy shape can also, under the right conditions, act like an airplane wing and be lifted by the streamflow more readily than would a spherical particle of similar weight. Once this occurs for even a few particles, the imbrication is compromised and significant portions of the streambed become mobile.

Surficial Geology

Surficial geology is concerned with the material covering the bedrock. In the Catskills this surface material is principally soils and glacial deposits. The focus here is on a brief introduction to the glacial geology of the Schoharie watershed and stream corridor. The Greene County Soil Survey is an excellent source for examining the soils of the Upper Schoharie Creek corridor (USDA, 1993).

The ice ages of the last 1.6 million years (Pleistocene Epoch) have left the latest mark on the Catskill landscape. Vast continental ice sheets and smaller local mountain glaciers scoured the mountains and left thick deposits of scoured sediment in the valleys. The last ice sheet (the “Laurentide Ice Sheet”) reached maximum thickness over the Catskills about 22,000 years ago (Isachsen, et al., 2000) and had fully retreated by 12,000 years ago (Figure 2.5.2). As measured on the scale of geologic time this was a very recent event.



Figure 2.5.2. (a) Map of Laurentide ice sheet. (b) Photo of Greenland ice sheet in mountainous terrain.

The most recent ice ages – the time that spanned the last 30,000 years or so – had giant continental-sized ice sheets flowing across the northern landscape (Figure 2.5.2a). The ice sheet covering Greenland (Figure 2.5.2b) is a modern day analog to those Pleistocene conditions. The continental glaciers scoured and moved vast amounts of sediment across the landscape. Once the ice sheet started melting back into the Hudson-Mohawk River valleys to the north, smaller alpine glaciers formed in some of the higher mountains and further sculpted the landscape. The glaciers left a legacy that still profoundly influences hill slope and stream channel stability and water quality.

This was a period of accelerated erosion in the Catskills as the flowing ice sheet bulldozed sediment and “quarried” the bedrock. Glacial erosion broke the rock down into an entrained mixture of fragments ranging in size from boulders to clay. This mixture of saturated sediment was carried along by the ice and deposited as till (unsorted assemblage of glacial sediment) or as stratified “drift” if the sediment was subsequently sorted by melt-water streams. These glacial deposits filled in deep river ravines that once drained the landscape prior to the last glacier’s advance over the mountains.

As the climate warmed and ice thinned, the landscape was deglaciated – lobes of the continental ice sheet melted back from the central Catskills in periodic stages (Dineen, 1986). As the ice sheet pulled back (and occasionally re-advanced as distinct “lobes” of flowing ice) alpine glaciers formed on some of the newly exposed peaks (e.g. Hunter Mountain). Meltwater from the decaying ice left a



Figure 2.5.3. Example of extent of pro-glacial lake in adjacent Esopus Creek watershed

complex array of stream (outwash plain) and ice-contact (kame) sand and gravel deposits. Pro-glacial lakes formed where mountains, recessional moraines (deposits at former glacial margins) and ice impounded water and filled the valley floors with thick deposits of layered silt and clay (Figure 2.5.3). The extent of the pro-glacial lakes in the Catskills are inferred from elevations of “fossil” deltas from meltwater streams pouring into valley-filling. One long-standing lake during this time filled the ice-free parts of the Schoharie valley at an elevation up to 1600 ft (Rich, 1935) corresponding to the elevation of the notch at Grand Gorge. The notch was a spillway for Glacial Lake Grand Gorge, discharging water into the Delaware basin. The extent of the glacial lakes during the prolonged melting of stagnant continental ice exposed a large proportion of the catchment to the accumulation of layered fine sediment. As climate fluctuated during the period of deglaciation, temporary re-

advances of ice from ice sheet lobes or alpine glaciers would leave till and other meltwater deposits on top of the earlier glacial material, resulting in the complex lateral and vertical distribution of glacial deposits observed today. After the ice fully retreated north, rainfall-runoff returned as the predominant sculptor of the landscape.

Glacial geology sets the geologic framework for most of the Upper Schoharie Creek stream system, controlling such characteristics as depth of *alluvium* (water worked sediments), presence of non-alluvial boundary conditions (till and glacial lake sediments), sediment supply and stream channel slope and geometry. For example, glacial depositional features that partially fill river valleys, such as recessional moraines or kame terraces along the valley wall, influence valley slope and cause valley constriction, both of which limit where the river channel can occur. Also, locally complex stratigraphy of glacial till, glacial lake deposits and unconsolidated *fluvial* deposits in the stream bank profile significantly influence erosional processes. Understanding detailed glacial geology can help identify causes of stream erosion and water quality problems as well as assist in prioritizing where future stream stabilization or restoration actions may be most useful.

For more detail on the glacial geology of the Catskills the reader is referred to Rich (1935), Cadwell (1986), Dineen (1986) and for a popularized account Titus (1996). Figure 2.5.4 presents a map of the surficial geology for the Schoharie basin as mapped by Cadwell (1987). It is safe to say that the actual geology is significantly more complicated than depicted on such a small scale map.




Figure 2.5.4. Surficial Geology of the Schoharie Creek Basin as mapped at 1:250,000

Hydrogeology

Though groundwater is not the subject of this Management Plan, its constructive role in maintaining base flow to the stream and cold water springs for thermal refugia, and its destructive role in hill slope failures should be addressed.

Given that much of the valley floor stratigraphy includes buried impermeable layers of glacial lake silt and clay and/or glacial till, groundwater circulating through the upper permeable coarse-grained alluvium is often perched and discharges as springs or base flow to the stream. Following periods of excess rainfall not only does the stream flow increase to or near flood stage, but the water table also increases and can flood basements. Much of the “flood” damage to basements in the Catskills is due to excess groundwater in these shallow groundwater systems and not directly from stream flooding.

Groundwater flow through the complex glacial stratigraphy on the hill slopes is a major factor in the massive hill slope failures that impact stream channel conditions and water quality (Photo 2.5.5). The combination of stream erosion at the toe of the hill slope, fluctuating groundwater levels, differential seepage from the slopes and saturated sediment can result in very



Photo 2.5.5. Hillslope failure in the Schoharie basin

long-lasting, deep-seated slope failures. Examples abound throughout the watershed. Every major rainfall-runoff event seems to generate new slope failures or reactivate older failures. Some of the chronic turbidity sources in the tributary streams are from these hill slope failures, such as in Batavia Kill and West Kill (GCSWCD, 2003; GCSWCD, 2005).

Stream Channel Geology

Developing an effective stream corridor management plan that incorporates geologic boundary conditions requires an additional step beyond describing the geologic setting. Additional analysis is needed to characterize the surficial geology that forms the stream channel boundary by some of its sedimentologic conditions, specifically grain size distribution, cohesiveness, and consolidation.

Upper Schoharie Creek and its tributaries flow across a landscape characterized by sedimentological heterogeneity as a result of the complex distribution of glacial deposits and

landforms. Stream channel stability and water quality vary in part as a function of this heterogeneity. By classifying the surficial geology along the stream corridor into mappable units that describe the potential for bed and bank erosion and entrainment of the stream channel material, recommendations for management of stream reaches can better reflect local geological considerations.

Rubin (1996) began this effort in the Stony Clove basin by classifying the glacial deposits into three sedimentologic units and mapping their distribution along the Stony Clove main stem and tributary channels (GCSWCD, 2004). The following are the three key sedimentologic units that influence water quality and stream stability. They were first proposed by Rubin (1996), and have been subsequently adapted for the development of stream management plans (GCSWCD, 2004; GCSWCD, 2005; CCE, 2007).

Unconsolidated Deposits

This general term is applied to all unconsolidated deposits regardless of whether they were deposited directly as post-glacial stream deposits, glacial outwash (proglacial fluvial sediments), reworked outwash, *kame terrace* deposits, melt-out till, *moraine* deposits or reworked lodgement till (Photo 2.5.6). The unit is composed of sand, gravel, cobbles, boulders and a



Photo 2.5.6. Coarse fluvial sediment comprises most of the Schoharie Creek and East Kill stream banks

small clay/silt fraction. The unconsolidated deposits are present in valley centers, typically ranging from four to twelve feet in thickness (Rubin, 1996). With the exception of a thin, weathered mantle often capping it, this is the uppermost geologic unit most commonly forming stream banks. Boulders specific to this geologic unit naturally drop out as stream banks are eroded, providing some aquatic habitat and diversity.

Lacustrine silt/clay

Reddish or pinkish brown, finely-layered, silty-clay deposits are present in significant portions of the Upper Schoharie Watershed (Photo 2.5.7). It was deposited *subaqueously* (from streams discharging into one or more glacial lakes) as a sediment blanket draped over underlying till or bedrock. Locally, it was also deposited in smaller impoundments



Photo 2.5.7. Clay deposit along the Schoharie Creek

associated with alpine glaciers and moraine dams. It is commonly exposed along the toe of the stream bank, sometimes in the channel bottom (often beneath a thin cover of coarse alluvium), and less frequently as long and/or large banks.

The fine, uniform grain size results in a very cohesive deposit that exhibits unique hydraulic and mechanical erosion characteristics. While the silts are easily entrained under high runoff events, many of the clay-rich deposits are resistant to hydraulic erosion. Susceptibility to erosion is largely dependent upon whether the layered silt/clay has been mechanically disturbed by geotechnical failures or human disturbance. The silt/clay unit tends to erode mechanically by slumping along rotational faults, subsequently losing its layered structure and cohesive strength (Figure 2.5.5). Within the silt and clay layers, strata of sand sometimes occur, creating the potential for piping and associated mechanical failures. When saturated, it tends to be extremely soft and in this physically- and chemically-weakened condition is susceptible to creep and erosion. Research in the Esopus Creek also demonstrated that erodibility was a function of the degree of disturbance of the exposure surface, and tended to diminish over time as the exposed surface smoothed through erosion (Fischenich et al., 2007).

Glacial Lake Clays and Stream Bank Erosion

Stream banks formed in deep clay deposits tend to fail by rotational failure which occurs in cohesive materials when a block of disturbed bank material slides along a curved failure surface (fault). The block tends to rotate (appears to “slump”) back toward the bank as it slides, in a rotational slip.




Figure 2.5.5. Glacial lake clays and stream bank erosion

Where vegetative cover is lost and large exposures of lacustrine silt/clays occur, revegetation is usually slow to due to the poor drainage and rooting characteristics of the soil. A metal probe or stick can often be sunk into this unit to depths of between three and five feet, thus enabling identification even when it is covered by a thin cobble layer. Elongate troughs, scour holes and even deep potholes reflect its entrainment potential during scouring flows. Clear stream water contacting lake clays often results in an entire stream becoming turbid within 50 feet. In the Upper Schoharie Creek watershed this lacustrine silt/clay, along with lodgement till, are the primary sources for suspended sediment and turbidity problems.

Lodgement Till

Lodgement till is an over-consolidated (very dense), clay-rich, reddish brown deposit that is prevalent in the Upper Schoharie Creek and East Kill watersheds (Photo 2.5.8). This hard-packed silty clay with embedded pebbles, cobbles and boulders forms a number of steep banks in the drainage



Photo 2.5.8. Lodgement till in the East Kill

basin. Its dense, consolidated character is distinguished from the looser assemblage of mixed sediment sizes (silty sand-boulder) that comprises melt-out till found in moraines and along mountain sides. Lodgement till is typically exposed in stream channels where overlying lake clay deposits have been removed by erosion, where streams have scoured into valley wall deposits or where they have breached morainal ridges.

Its relatively competent nature, especially compared to disturbed lacustrine sediment, make it significantly more resistant to hydraulic erosion. It is however, susceptible to mechanical erosion by mass failure of fracture bound blocks during saturation/desaturation and freeze/thaw cycles. This failed material is subsequently eroded by streamflows. Under conditions of high stream velocities and discharges, lodgement till is a contributor of sediment. However, where the stream (particularly in tributary valleys) is against the valley wall and the hill slope composed of lodgment till is saturated, long-lasting exposures can be chronic sources of suspended sediment into the stream well-after a storm event. Reaches in many of the Schoharie tributaries are subjected to this phenomenon. Rain water and overland runoff contacting exposed banks can also readily entrain sediment from these units. For field mapping, a metal probe or stick can rarely be pushed into this unit more than 0.2 feet.

Bedrock Control

The presence of bedrock sills and banks is an additional geologic unit equally important in characterizing geology for stream corridor management. These hydraulic controls can represent natural limits to changes in the stream channel system caused by incision or lateral migration. Examples include the falls in the headwater reaches, and occasional bedrock stream banks and sills along the Upper Schoharie Creek/East Kill (Photo 2.5.9).



Photo 2.5.9. Lateral bedrock control in the East Kill

In summary, the variable character of the Upper Schoharie watershed is largely a reflection of the geologic bedrock control and complex glacial history of the valley. These geologic influences are evident in the sedimentological variation characterizing the topography and geomorphology of the stream channel boundary. The nature of these deposits makes them variably susceptible to stream erosion. In particular, the lacustrine and till sediments are sensitive to natural or man made disturbances which can have a long lasting negative effect on channel stability, water quality and stream ecology.

Stream Management Implications

The inclusion of geology in stream management consideration for Upper Schoharie Creek and East Kill generally falls into four categories: fluvial erosion, hill slope erosion, water quality, and sediment supply.

Fluvial erosion

There are different types or “styles” of stream bank erosion associated with the different geologic units the stream encounters. The prediction, prevention and/or treatment of the eroding stream bank must factor in the stream bank material composition and the underlying mechanism of failure. Observations made during this planning process and previous similar projects throughout the watershed indicate the following:

- Pro-glacial lake sediment erodes easily during storm events once exposed; however, if the “soft” silt and clay unit is overlain by coarser fluvial sediment (sand-boulder sized material) it is typically a short-lived exposure and the stream bank tends to get armored by the draping of the coarser sediment.
- Pro-glacial lake deposits that are undisturbed are much more resistant to erosion than those that have had their physical and chemical bonds weakened by mechanical action (including abrasion and displacement from hill slope failures).
- Glacial till tends to erode either as (a) mass slumping from saturated conditions or (b) translational fracture-bound failures forming high steep banks.

- Coarse-grained, non-cohesive fluvial sediment will erode easily if not protected by dense roots or revetment.

Hill slope erosion

The mass wasting, or geotechnical failure of the valley hill sides when proximal to stream channels can result in chronic and excess fine and coarse sediment supply. This is a relatively common problem in the tributary valleys. Sediment entrainment occurs as a result of exposed glacial till or disturbed lake deposits to flood flows. In extreme situations, debris flows from these failures may block or cause the stream channel to adjust its planform. If the adjacent hill slope erosion is from a geotechnical failure in glacial till or pro-glacial lake sediment and the stream is actively eroding into the toe of the hill slope the problem is perpetuated by constantly activating the failure. Stream restoration or road construction/repair in these settings must first address whether the geotechnical failure can be resolved before dealing with the stream channel stabilization. Future construction or development activities in the Schoharie Creek tributary valleys should include geotechnical investigations and slope stability analyses to ensure that the proposed actions do not contribute to new slope failures or exacerbate existing failures.

Water quality

The “muddy” or turbid water that follows a storm event carries the fine silt and clay particles initially deposited as glacial till or pro-glacial lake sediment (Photo 2.5.10). Fluvial and hill slope erosion of these fine sediment sources, along with re-suspension of fine sediment deposited in the stream bed are the primary cause of the turbid water conditions. The



Photo 2.5.10. Turbid tributary entering the Schoharie Creek during a summer storm.

fact that the glacial till and glacial lake sediment is widely distributed throughout most of the watershed suggests that effective removal of the stream from contacting this material

is impractical to consider. High levels of suspended sediment and associated turbidity have been and will be an ongoing water quality condition in the Upper Creek watershed.

Sediment supply

The mantle of glacial deposits over the landscape is the primary source material for all the coarse and fine sediment that the stream system conveys. At any given time along any given reach of stream most of the sediment observed has been in the stream system for a “long time”. However, it is important to determine where sediment recruitment takes place. Unanswered questions remain: Which tributary streams deliver a proportionally larger amount of bed load material that Schoharie Creek has to process? Are there localized sources in the watershed that lead to localized aggradation?

Recommendations

The following recommendations are presented as an initial scope for further investigation and development of products to improve the Upper Schoharie Creek.

- Work with research and/or academic institutions to better characterize the lateral and vertical distribution of glacial deposits that influence stream channel condition and water quality. Encourage academic interest in addressing this applied geology issue.
- Continue to monitor previously mapped fine sediment sources along Upper Schoharie Creek and East Kill, and implement a program to identify “new” exposures. The aim of this effort is to better characterize the temporal nature of fine sediment exposures and their contribution to water quality problems in the basin.
- Using (1) georeferenced data obtained during the geomorphic investigation, (2) available soils map and (3) further reconnaissance mapping develop a stream channel geologic map for Upper Schoharie Creek and East Kill.
- Extend stream channel geologic and fine sediment source mapping into all tributary valleys not previously assessed, and develop a sediment budget to include more detail on the tributaries so that the relative contribution of

sediments from these sources can be determined and the potential benefits of management actions in the tributaries better elucidated.

- Support an investigation of the geotechnical and hydrogeologic processes controlling coupled hill slope and stream bank erosion in order to evaluate management feasibility.
- Develop a document that informs stream managers how to use this information when designing and implementing stream stabilization projects in the region.

References

- Cadwell, D.H. 1986. Late Wisconsinan stratigraphy of the Catskill Mountains. in *The Wisconsinan Stage of the First Geological District, Eastern New York*, ed. Donald H. Cadwell, NYS Museum Bulletin 455: 192 p.
- Cadwell, D.H. Surficial Geologic Map of New York 1:250,000: Hudson-Mohawk Sheet, 1987; Map and Chart Series No. 40. NYS Museum.
- CCEUC. 2007. Upper Esopus Creek Management Plan. Cornell Cooperative Extension of Ulster County, Kingston, NY.
- Dineen, R.J. 1986. Deglaciation of the Hudson valley between Hyde Park and Albany, NY in *The Wisconsinan Stage of the First Geological District, Eastern New York*, ed.
- Fischenich, J.C., Channell, M.G. and Davis, D. 2007. Stability of Fine-grained Glaciogenic Sediments in the Upper Esopus Creek Corridor, NY. In: *Upper Esopus Creek Management Plan, Volume III, Appendix D*, Cornell Cooperative Extension of Ulster County, Kingston, NY.
- Fisher, D.W., Y.W. Isachsen, and L.V. Rickard. 1970. Geologic Map of New York State: Hudson-Mohawk Sheet. NYS Museum.
- GCSWCD, 2003. Batavia Kill Stream Management Plan. Greene County Soil and Water Conservation District, Cairo, NY.
- GCSWCD, 2004. The Stony Clove Stream Management Plan. Greene County Soil and Water Conservation District, Cairo, NY.
- GCSWCD, 2005. West Kill Stream Management Plan. Greene County Soil and Water Conservation District, Cairo, NY.
- Isachsen, Y.W. et al., editors. 2000. *Geology of New York: A simplified account*, 2nd ed. NYS Museum Educational Leaflet 28: 294 p.

- Rich, J.L. 1935. Glacial geology of the Catskills. NYS Museum Bulletin 299, 180 p.
- Rubin, P. 1996. Geologic Mapping of Sediment Sources. Draft NYCDEP memorandum.
- Schneiderman, J.S. 2003. The Earth Around Us: Maintaining a Livable Planet. Boulder: Westview Press.
- Titus, R. 1998. The Catskills: A geological guide. Purple Mountain Press. Flesichmanns, NY.
- USDA. 1993. Soil Survey of Greene County, New York. United States Department of Agriculture, Soil Conservation Service. Available from GCSWCD (www.gcswcd.com), Cairo, NY.
- VanAller Hernick, L. 1996. The Gilboa Fossils. Givetian Press, Rensselaerville, NY

2.6 Wetlands & Floodplains

Wetlands are areas where the soil is inundated by surface or ground water often enough that the prevalent vegetation community is adapted for life in saturated soils. The term “wetlands” covers a diverse set of conditions, including swamps, marshes, bogs and fens. The timing and duration of soil saturation largely determines how the soil develops and the particular community of plants and animals living in and on the soil. The prolonged presence of water creates conditions that favor the growth of specially adapted plants (hydrophytes) and promote the development of characteristic wetland (hydric) soils (USEPA, 2006).

Wetlands are recognized as important features in the landscape that provide numerous beneficial functions, including protecting and improving water quality, providing fish and wildlife habitats, storing floodwaters, and maintaining surface water flow during dry periods. These valuable services are the result of the inherent and unique natural characteristics of wetlands (USEPA, 2001).

It has long been accepted that the presence and position of wetlands play a key role in the filtration of dissolved inorganic nutrients and suspended materials from water (Johnston et al. 1990; Welsh et al., 1995; Hammer, 1997). The wetland filtration function buffers our streams, ponds, and lakes from receiving excess nutrients and suspended materials. The addition of excessive amounts of nitrogen and phosphorous to marine and freshwater systems, where nitrogen and phosphorous, respectively, are generally limiting to plant growth, can lead to accelerated eutrophication (USEPA Office of Water, 1997).

Nitrate, the most mobile form of nitrogen, can either be absorbed by vegetation, leach into groundwater or surface water, or be converted to nitrogen gas in the process of denitrification (Welsh et al., 1995). Sediment and phosphorous retention is also an important function of wetland systems, with excess phosphorous tending to be associated with sediment. Wetland trapping and storage of sediment will also trap and store excess phosphorous. However, sediment attached phosphorous is subject to resuspension and movement when wetland sediments are disturbed (Welsh et al., 1995). Fecal coliform bacteria are often associated with suspended materials as well. Many of the organisms

associated with fecal coliform bacteria cannot survive for long periods of time outside of their host organism. Therefore, the wetland function of retaining suspended material promotes the die-off of the fecal coliform bacteria (Johnston et al., 1990). This function is threatened with increasing storm water runoff and loss of wetlands. As of the mid 1980's, New York State had lost 60% of its original wetlands (Mitsch, 1993).

Fertilizer application, septic systems, and sewage treatment plant discharges directly affect nutrient exports (Caraco and Cole, 1999; Vitousek et al., 1997). Clearing of land for building lots and agriculture is often associated with a decrease in wetland acreage. The loss of wetlands removes a significant sink for fixed nitrogen leading to an increase in the mobility of nitrogen to streams, rivers, and lakes (Vitousek et al., 1997). Wetland restoration may be the most cost-effective method of decreasing nitrogen pollution (Carpenter et al., 1998). Removal of wetland systems has similar effects on sediment and phosphorous exports.

Federally Designated Wetlands

The National Wetlands Inventory (NWI) of the U.S. Fish & Wildlife Service produces information on the characteristics, extent, and status of the Nation's wetlands and deepwater habitats (USFWS, 2003). According to NWI maps there are 168 federally designated wetlands totaling 311 acres, including open water in the East Kill Watershed. These wetlands are 84.4% Palustrine, 9.3% Lacustrine, 6% Riverine, and 0.2% Upland.

The Palustrine System includes all nontidal wetlands dominated by trees, shrubs, emergents, mosses or lichens, and all such wetlands that occur in tidal areas where salinity due to ocean derived salts is below 0.5 ppt. Wetlands lacking such vegetation are also included if they exhibit all of the following characteristics: are less than 8 hectares (20 acres); do not have an active wave-formed or bedrock shoreline feature; have at low water a depth less than 2 meters (6.6 feet) in the deepest part of the basin; and have a salinity due to ocean-derived salts of less than 0.5 ppt. The Lacustrine System includes wetlands and deep water habitats situated in a topographic depression or a dammed river channel, lacking trees, shrubs, persistent emergent vegetation, emergent mosses or lichens with greater than 30% areal coverage, a total area exceeding 8 hectares (20 acres). The Riverine System includes all wetlands and deep water habitats contained in natural or artificial channels periodically or

continuously containing flowing water or which forms a connecting link between the two bodies of standing water. Upland islands or Palustrine wetlands may occur in the channel, but they are not part of the Riverine System. Upland systems includes all areas not defined as wetland or deep water habitats. (USFWS₁, 2006).

In 2003, the dominant wetland type in the East Kill Watershed was Palustrine emergent wetlands (42.8%) (Table 2.6.1). Palustrine wetlands are vegetated wetlands including the small, shallow, permanent or intermittent water bodies often called ponds. Palustrine wetlands may be situated shoreward of lakes, river channels, or estuaries; on river floodplains; in isolated catchments; or on slopes. They may also occur as islands in lakes or rivers. Palustrine emergent wetlands are characterized by erect, rooted, herbaceous hydrophytes, excluding mosses and lichens. This vegetation is present for most of the growing season in most years. These wetlands are usually dominated by perennial plants. The remaining wetlands in the watershed are dominated by of four major types: Palustrine unconsolidated bottom (18.5%), Palustrine forested (12.5%), Palustrine scrub-shrub (10.7%), and Lacustrine lemnetic unconsolidated bottom (9.3%). These wetlands are scattered throughout the watershed.

NWI Code	NWI Wetland Classification	Acres	%
L1UB	Lacustrine, Limnetic, Unconsolidated Bottom	28.9	9.3
PEM	Palustrine, Emergent	133.0	42.8
PFO1A	Palustrine, Forested	38.8	12.5
PSS1A	Palustrine, Scrub-Shrub	33.2	10.7
PUBFb	Palustrine, Unconsolidated Bottom	57.4	18.5
R3	Riverine, Upper Perennial	18.7	6.0
U	Upland	0.7	0.2

Palustrine unconsolidated bottom wetlands include all wetlands and deep water habitats with at least 25% cover of particles smaller than stones (less than 6-7 cm), and a vegetative cover less than 30%. Palustrine forested wetlands are characterized by woody vegetation that is 6 m tall or taller. Palustrine scrub-shrub wetlands include areas dominated

by woody vegetation less than 6m (20 feet) tall. The species included in this wetland type are true shrubs, young trees (saplings), and trees or shrubs that are small or stunted because of environmental conditions. Lacustrine limnetic unconsolidated bottom wetlands are situated in a topographic depression or a dammed river, are deep-water habitats (>6.6 feet deep) lacking vegetation over 30% of its area and must exceed 20 acres in size. These relatively large, deep water habitats have bottoms with more than 25% of their particles smaller than stones (<6-7 cm) (USFWS₁, 2006). In the East Kill Watershed, Colgate Lake accounts for the entire area of this wetland type.

Federally designated wetlands are protected under the Clean Water Act, a 1977 amendment to the Federal Water Pollution Control Act of 1972, which set the basic structure for regulating discharges of pollutants to waters of the United States (USEPA₁, 2003). Section 404 of the Clean Water Act established a program to regulate the discharge of dredged and fill materials into waters of the United States, including wetlands. Activities in waters of the United States that are regulated under this program include fills for development, water resource projects (such as dams and levees), infrastructure development (such as highways and airports), and conversion of wetlands to uplands for farming and forestry (USEPA, 2003).

New York State Designated Wetlands

The Freshwater Wetlands Act (FWA), Article 24 of the Environmental Conservation Law, provides NYS DEC and the Adirondack Park Agency (APA) with the authority to regulate freshwater wetlands in the state. The NYS Legislature passed the Freshwater Wetlands Act in 1975 in response to uncontrolled losses of wetlands and problems resulting from those losses, such as increased flooding. The FWA contains the following Declaration of Policy:

"It is declared to be the public policy of the state to preserve, protect and conserve freshwater wetlands and the benefits derived therefrom, to prevent the despoliation and destruction of freshwater wetlands, and to regulate use and development of such wetlands to secure the natural benefits of freshwater wetland, consistent with the general welfare and beneficial economic, social, and agricultural development of the state (ECL Article 24-0103)."

The FWA protects those wetlands larger than 12.4 acres (5 hectares) in size, and certain smaller wetlands of unusual local importance. The law requires DEC and APA to map those wetlands that are protected by the FWA. In addition, the law requires DEC and APA to classify wetlands. Outside the Adirondack Park, DEC classifies wetlands according to 6NYCRR Part 664, Wetlands Mapping and Classification Regulations from Class 1, wetlands which provide the most benefits, to Class IV, wetlands which provide the fewest benefits. Around every regulated wetland is a regulated adjacent area of 100 ft., which serves as a buffer area for the wetland (NYS DEC, 2003).

According to DEC maps, there are 7 NYSDEC designated wetlands totaling 119 acres, within the East Kill watershed. All of these wetlands are designated Class 2. These wetlands are scattered throughout the watershed.

Both Federal and NYS Designated Wetlands maps are available at County Soil & Water Conservation District Offices. It must be cautioned that these maps should only be used as guidance of wetland locations and boundaries. It is the responsibility of property owners to determine if wetland areas will be disturbed by proposed projects. Smaller wetlands which meet federal criteria may not have been mapped but are still protected by federal regulations. The NYSDEC offers wetland delineation services to landowners when they need more precise information, such as when they are planning to conduct work near a NYSDEC designated wetland area.

Floodplains

A floodplain is streamside land that gets periodically inundated by floodwaters. Floodplains are important because they temporarily store floodwaters, improve water quality, and provide important habitat for wildlife. Natural floodplains help reduce the heights of floods. During periods of high water, floodplains serve as natural sponges, storing and slowly releasing floodwaters. The floodplain provides additional "storage," reducing the velocity of the river and increasing the capacity of the river channel to move floodwaters downstream. Natural floodplains also help improve water quality. As water courses through the floodplain, plants serve as natural filters, trapping sediments and capturing pollutants (American Rivers, 2003).

One of the largest problems facing floodplain management is the disconnection of a stream from its floodplain. Management practices such as channelization, straightening, development, and loss of riparian vegetation may lead to stream channel *incision* or down-cutting. As the stream incises it will lower the streambed elevation, no longer allowing floodwaters to spill out into the floodplain. As a result flood velocity will increase causing streambank *degradation* until a new floodplain is created at the lower streambed elevation. Building homes within the floodplain is incompatible with proper floodplain function. Many people want to live by streams but as they develop the floodplain, they often increase stream degradation by undertaking stream management activities to protect their property from flooding.

The Federal Emergency Management Agency (FEMA) performs hydrologic and hydraulic studies to produce Flood Insurance Rate Maps (FIRM), which identify flood-prone areas (FEMA, 2003). These studies analyze the data from local streamflow *gages* to predict how frequently different floods will occur, and to determine the magnitude of the benchmark “100-year flood”. This is the flow that has a statistical probability of recurring once every 100 years, but because it is a statistical prediction, based on historical record, “100-year floods” could be seen more or less frequently than every hundred years, especially if changes in climate or land use occur. An engineering model is then used to map the predicted boundaries of the 100-year flood on the floodplain. Towns then use these maps to help determine areas where the risk of flooding is high enough to warrant special precautions or review of land development. Towns are required to pass a floodplain protection ordinance that sets certain limits on building in the 100-year floodplain in order to participate in the National Flood Insurance Program.

Some towns develop other ordinances that help focus review of development on lands that could affect stream and floodplain function. One example of an innovative model in effect locally is the ordinance that was adopted by the Town of Woodstock. The text of this ordinance can be found at <http://www.woodstockny.org/Laws/WWLAW9.pdf>

Digital Flood Mapping Project

The NYS DEC Bureau of Program Resources and Flood Protection has developed new digitized floodplain maps, using topographic information derived from an airborne laser imaging technology called LIDAR (Light Detection and Ranging). LIDAR data, together with updated computer HEC models and digital aerial photography, enable engineers to produce extremely detailed and accurate maps. Modeling with this new data allows for flood contour lines indicating various depths of water under 100-year and other flood conditions. FEMA's new hardcopy Flood Insurance Rate Maps (FIRMs) are a vast improvement over their predecessors. One of the most obvious improvements is the inclusion of base map imagery utilizing the 2004 orthoimagery from New York's statewide orthoimagery program. A New York State Floodplain Management Map (NYSFMM) series has also been developed to provide floodplain managers, municipal planners, and other professionals with a tool for mitigation and planning. In addition to the information found on a FIRM, the NYSFMMs also contain department-set survey reference marks and flood depth contours (NYSDEC, 2006).

The new FIRM hardcopy maps are available for viewing at County Soil & Water Conservation District Offices and most town halls. Using GIS mapping software, Greene County Soil & Water Conservation District (GCSWCD) is able to overlay tax parcel boundaries with digital floodplain boundaries to assess if a property falls within a flood zone. This service is available to all interested. Floodplain maps of each management unit can be found in Section 4 Management Unit Summary & Recommendations.

References

- American Rivers. 2003. The value of floods and floodplains. Available of web: <http://www.amrivers.org/index.php?module=HyperContent&func=display&cid=1823>.
- Caraco, N.F., and J.J. Cole 1999. Human Impact on Nitrate Export: An Analysis Using Major World Rivers. *Ambio* 28(2):167-170.
- Carpenter, S.R., Caraco, N.F., Correll, D.L., Howarth, R.W., Sharpley, A.N., and Smith, V.H. 1998. Nonpoint Pollution of Surface Waters With Phosphorous and Nitrogen. *Ecological Applications* 8(3): 559-568.
- Dahl, T.E. 2006. Status and Trends of Wetlands in the Conterminous United States 1998 to 2004. U.S. Department of the Interior; Fish and Wildlife Service, Washington, DC. 112 pp.
- FEMA. 2003. FEMA, Map Service Center. Federal Emergency Management Agency. Available on web: www.msc.fema.gov/prodover.shtml.
- Hammer, Donald A. 1997. Creating Freshwater Wetlands, second edition. Lewis Publishers, NY.: 106-116 P.

- Johnston, Carol A., Detenbeck, Naomi E., and Niemi, Gerald J. 1990. The cumulative effect of wetlands on stream water quality and quantity. A landscape approach. *Biogeochemistry* 10: 105-142.
- Mitsch, Willaim J. and Gosselink, James G. 1993. *Wetlands*, second edition. Van Nostrand Reinhold, NY: 46-47 P.
- NYSDEC. 2003. Programs to Conserve Wetlands. Available on web: <http://www.dec.state.ny.us/website/dfwmr/habitat/fwwprog4.htm>.
- NYSDEC. 2006. NYS DEC Floodplain Mapping Program Overview. Available of web: <http://www.dec.state.ny.us/website/dow/wgit/fpm/index.html>.
- USEPA Office of Water. 1997. Monitoring Guidance for Determining the Effectiveness of Nonpoint Source Controls. EPA-841-B-96-004. United States Environmental Protection Agency, Washington, D.C.: 1-4 P.
- USEPA. 2001. Functions and Values of Wetlands. United States Environmental Protection Agency, Washington, D.C. Publication 843-f-01-002c. Available on web: http://www.epa.gov/owow/wetlands/pdf/fun_val.pdf.
- USEPA, 2003. Section 404 of the Clean Water Act: An Overview. United States Environmental Protection Agency, Washington, D.C. Available on web: <http://www.epa.gov/owow/wetlands/facts/fact10.html>.
- USEPA₁. 2003. Wetland Laws of the United States. United States Environmental Protection Agency, Washington, D.C. Available on web: <http://www.epa.gov/owow/wetlands/laws/>
- USEPA. 2006. What are Wetlands? United States Environmental Protection Agency, Washington, D.C. Available on web: <http://www.epa.gov/owow/wetlands/vital/what.html>.
- USFWS. 2006. Information about National Wetland Inventory. US Fish & Wildlife Service, Division of Habitat and Resources Conservation, Arlington, VA. Available on web: <http://www.nwi.fws.gov/aboutus.htm>.
- USFWS₁. 2006. Wetlands and Deepwater Habitats Classification National Wetlands Inventory Mapping Code Description. US Fish & Wildlife Service, Division of Habitat and Resources Conservation, Arlington, VA. Available on web: <http://wetlandfws.er.usgs.gov/NWI/webatx/atx.html>.
- Vitousek, Peter M., Aber, John D., Howarth, Robert W., Likens, Gene E., Matson, Pamela A., Schindler, David W., Schlesinger, William H., and Tilman, David G. 1997. Human Alteration of the Global Nitrogen Cycle: Sources and Consequences. *Ecological Applications* 7(3): 737-750.
- Welsch, David J., Smart, David L., Boyer, James N, Minkin, Paul, Smith, Howard C., and McCandless, Tamara L. 1995. Forested Wetlands, Functions, Benefits, and the Use of Best Management Practices. United States Department of Agriculture Forest Service, Radnor, PA.: 30-31 P.

2.7 Riparian Vegetation Issues in Stream Management

General Concepts of Riparian Vegetation Ecology and Management

The Role of Vegetation in Maintaining a Healthy Stream

Although people value trees and plants along a stream for their contribution to the beauty of the streamside landscape, the vegetation in a watershed, especially in the *riparian* area, plays a critical role in providing for a healthy stream system. The riparian, or streamside, plant community serves to maintain the riverine landscape and moderate conditions within the aquatic ecosystem.

As rainfall runs off the landscape, riparian vegetation:

- Slows the rate of runoff
- Captures excess nutrients carried from the land
- Protects stream banks and floodplains from the erosive force of water
- Regulates water temperature changes

It also:

- Provides food and cover to terrestrial and aquatic fauna
- Conserves soil moisture, ground water and atmospheric humidity.

Vegetation's Erosion and Pollution Prevention Capabilities

Riparian vegetation serves as a buffer for the stream against activities on upland areas. Most human activities whether agriculture, development, or even recreation, can result in a disturbance or *discharge* which can negatively impact the unprotected stream. Riparian vegetation captures and stores pollutants in overland flow from upland sources such as salts from roadways and excess fertilizer from lawns and cropland. The width, density, and structure of the riparian vegetation community are important characteristics of the buffer that can be used to define the level of its functionality.

On bare soils, high stream flows can result in bank erosion and overbank flow can cause soil erosion and scour on the floodplain. The roots of vegetation along the bank hold the soil and shield against erosive flows. On the floodplain, vegetation slows flood flows, reducing the energy of water. This reduction in energy will decrease the ability of water to

cause erosion and scour. Furthermore, as vegetation slows the water, the soil suspended in the water is deposited on the floodplain.

Vegetation's Hydrologic Influences

Vegetation intercepts rainfall and slows runoff. This delay increases the amount of precipitation that infiltrates the soil and reduces overland runoff. A reduction and delay in runoff decreases the occurrence of destructive flash floods, lowers the height of flood waters, and extends the duration of the runoff event. These benefits are generally most readily observed in forested watersheds such as the East Kill, as opposed to similar watersheds where urban development is the chief land use (Figure 2.7.1). The reduction in flood stage and duration typically means fewer disturbances to stream banks and floodplains.




Figure 2.7.1. Comparison of Runoff on a Forested Watershed Versus a Deforested Watershed. Illustration by P. Eskeli 2002, from Watershed Hydrology, P.E. Black, 1991, Prentice Hall, page 202, 214.

Ecological Importance of Vegetation in the Riparian Zone

Vegetation along a stream also functions to provide the climate, habitat, and nutrients necessary for aquatic and terrestrial wildlife. Trees shading a stream help maintain cool water temperatures needed by native fish populations. Low hanging tree branches and roots on undercut banks create cover for fish from predators such as birds and raccoons. Natural additions of organic leaf and woody material provide a food resource needed by terrestrial insects and aquatic macroinvertebrates (stoneflies, mayflies, etc.), the primary source of food for fish. Terrestrial wildlife depends upon vegetation for cover as they move from the upland community to the water's edge. A diverse plant community, one similar to the native vegetation of the East Kill, provides a wide range of conditions and materials needed to support a diverse community of wildlife. If the vegetation is continuous within the riparian zone along the length of a stream, a corridor is available for wildlife migration. Connectivity between the riparian and upland plant communities enhances the ability of upland and riparian plant and animal communities to thrive despite natural or human induced stress on either community.

Characteristics of a Healthy Riparian Plant Community

A healthy riparian plant community should be diverse. It should have a wide variety of plants including trees, shrubs, grasses or herbs (Figure 2.7.2). The age of the plants should be varied and there should be sufficient regeneration of new plants to ensure the future of the community. A diverse community provides a multitude of resources and the ability to resist or recover from disturbance. An important difference between an upland plant community and a riparian community is that the riparian community must be adapted to frequent disturbance from flooding. Consequently, many riparian plants, such as willow, alder, or poplar can regrow from stump sprouts or can reestablish their root system if up-ended. Furthermore, the seed of these trees may have a greater ability to germinate and establish in depositional areas, such as gravel bars and lower flood benches.



Figure 2.7.2. A healthy riparian community is densely vegetated, has a diverse age structure and is composed of plants that can resist disturbance.

Riparian Vegetation of the East Kill

Forest History and Composition of the East Kill

Catskill mountain forests have evolved since the ice age reflecting the changes in climate, competition and human land use. The first of these changes was the result of the climatic warming that occurred after the ice age which enabled warm climate adapted plant communities to replace the cooler climate communities. Following the retreat of the glaciers, the forests of the East Kill basin gradually re-established and evolved from the boreal spruce/fir dominated forests, (examples of which can presently be found in Canada) to the maple-beech-birch northern hardwood forests (typical of the Adirondacks and northern New England) with the final transition of the lower elevations of the watershed to a southern hardwood forest dominated by oaks, hickory and ash (typical of the northern Appalachians). Dr. Michael Kudish provides an excellent documentation of evolution and site requirements of the region's forests in his book, The Catskill Forest: A History (Kurdish, 2000).

More recently, human activities have affected the forest either through the manipulation of regeneration for the maintenance of desirable species, the exploitation of the forest for wood and wood products or through development. Native American land management practices included the use of prescribed burning as a means of enabling the nut bearing oaks and hickories to remain dominant in the forest. European settlers contributing to the rising industrial economy in the 18th and 19th century greatly altered the landscape and

forest cover through land clearing for agriculture, forest harvesting for construction materials, and hemlock bark harvesting for the extraction of tannin. The land cover in the East Kill began to revert to forest with the local collapse of these economies in the 20th century and the acquisition of much of the land by the state for the Catskill Forest Preserve (Kudish, 2000). Please refer to individual Management Unit descriptions for more detailed information about past activities that affected the streamside and floodplain vegetation.

Previous land uses have had a significant role in determining the types of vegetation found along the stream. Due to the steepness of the sides of the valley, the most intensive development activities were confined to the valley floor along the stream. Pastures and fields were created from cleared, forested floodplains. After abandonment, these old fields have experienced a consistent pattern of recovery, with species dominating the initial regrowth including sumac, dogwoods, aspens, hawthorns, and white pine. These species are succeeded by other light loving hardwood tree species such as ash, basswood and elm or in lower parts of the watershed, hickories, butternut, and oak. Hemlocks are largely confined to the steeper stream banks and slopes where cultivation or harvesting of hemlocks for bark was impossible. More recent housing construction has re-intensified activity along the stream and been accompanied by the introduction of non-native vegetation typical of household lawns and gardens. Today, the East Kill watershed and riparian areas are largely forested, although the continuity of the forest is frequently interrupted by infrastructure, utility lines, residential land use and abandoned agricultural lands.

The Riparian Forest

Typically, a riparian forest community is composed of those plant species that thrive in a wet or moist location and have the ability to resist or recover from disturbance. Generally, the riparian forest community is more extensive where a floodplain or wetland exists and the side slopes to the valley are more gently sloping. The plant associations found in forested floodplain communities may be exclusive to riparian areas. Where the valley side slopes are steeper, the riparian community may occupy only a narrow corridor along the stream and then quickly transition to an upland forest community. From a vegetation assessment in a nearby watershed, it was found that northern hardwood communities on steep slopes adjacent to the stream contained a mix of ash, poplar, elm, beech, yellow birch and

some maple, whereas in upland northern hardwood communities, the yellow birch and maple became the dominant species. Soils, ground water and solar aspect may create conditions that allow the riparian forest species to occupy steeper slopes along the stream, as in the case where hemlock inhabits the steep, northfacing slopes along the watercourse.

Natural Disturbance and its Effects on the Riparian Vegetation

Riparian vegetation is disturbed by the forces of nature and development activities of those who live near the stream. Sources of natural disturbance include damage due to floods, ice floes, and to a lesser extent, high winds, pest and disease epidemics, drought and fire. Deer herds can also alter the composition and structure of the vegetation due to their specific browse preferences.

The 1996 flood created and reopened numerous high flow channels, reworked point bars, scoured floodplains and eroded formerly vegetated stream banks in the East Kill. Immediately following the flood, the channel and floodplains were scattered with woody debris and downed live trees. In the years since this event, much of the vegetation has recovered. Trees and shrubs flattened by the force of floodwaters have re-established their form. Gravel bars and sites disturbed in previous flood events became the seedbed for herbs and grasses. This type of natural regeneration is possible where the stream is stable and major flood events occur with sufficient interval to allow establishment. The effect of flood disturbance on vegetation along stable stream reaches is short term and the recovery/disturbance regime can be cyclical. If the disturbance of floods and ice are too frequent, large trees will not have the opportunity to establish. Typically, the limit that trees can encroach upon the channel is defined by the area disturbed by the runoff event that achieves bankfull flow (expected to occur on average every 1.3 years). While shrubs like willow and alder or herbaceous plants like sedges, which reestablish themselves quickly after disturbance, can grow in the bankfull channel, it is unadvisable to plant trees in this channel area.

Local geology and stream *geomorphology* may complicate the recovery process. A number of sites were found in the East Kill where vegetation has not been able to reestablish itself on the high, steep bank failures created during recent flood events. On these sites it will be necessary to understand the cause of the failure before deciding on whether to attempt

planting vegetation to aid in site recovery. In these instances, the hydraulics of the flowing water, the morphological evolution of the stream channel, the geology of the stream bank, and the requirements and capabilities of the vegetation must be considered before attempting restoration. Since the geologic setting on these sites is partially responsible for the disturbance, the period required for natural recovery of the site would be expected to be significantly longer unless facilitated by restoration efforts.

The ice break up in the spring, like floods, can damage the established vegetation along the stream banks and increase mortality of the young tree and shrub regeneration. Furthermore, ice floes can cause channel blockages which result in erosion and scour associated with high flow channels and overbank flow. Typically this type of disturbance also has a short recovery period.

Pests and diseases that attack vegetation can also affect changes in the ecology of the riparian area and could be considered a disturbance. The hemlock woolly adelgid (*Adelges tsugae*) is an insect, which feeds on the sap of hemlocks (*Tsuga spp.*) at the base of the



Hemlock woolly adelgid on the underside of a branch.

needles causing them to desiccate and the tree to take on a greyish color. Stress caused by this feeding can kill the tree in as little as 4 years or take up to 10 years where conditions enable the tree to tolerate the attack (McClure, 2001). This native insect of Japan was first found in the U.S. in Virginia in 1951 and has spread northward into the Catskills (Adams, 2002).

In the eastern United States, the adelgid attacks eastern hemlock (*Tsuga canadensis*) and Carolina hemlock (*Tsuga carolinianna* Engelman) and can affect entire stands of hemlock. Once a tree is infested, population trends for the insect are typified by a fluctuating density of the insect with some hemlock regrowth occurring in periods when population densities are lower. This regrowth is stunted and later attacked as the adelgid population increases. With each successive attack tree reserves become depleted and eventually regrowth does not occur. The native predators of hemlock woolly adelgid have not offered a sufficient biological control, but recent efforts to combat

the insect include experimentation with an Asian lady beetle (*Pseudoscymnus tsugae* Sasaji) which is known to feed on the adelgid. Initial experimental results have been positive, but large-scale control has yet to be attempted. The US Forest Service provides extensive information about this pest at its Morgantown office “forest health protection” webpage: www.fs.fed.us/na/morgantown/.

With respect to stream management, the loss of hemlocks along the banks of the East Kill poses a threat to bank stability and the aquatic habitat of the stream. Wildlife, such as deer and birds, find the dense hemlock cover to be an excellent shelter from weather extremes. Finally, dark green hemlock groves along the stream are quiet, peaceful places that are greatly valued by the people who live along the East Kill. The Olive Natural Heritage Society, Inc. is monitoring the advance of the hemlock woolly adelgid in the Catskills and is working in cooperation with NYS DEC on testing releases of *Pseudosymnus tsugae*. Initial results of the monitoring suggested a possible link between the presence of hemlock woolly adelgid on a site and the degree to which people use or access the site. Due to the widespread nature of the infestation, the use of chemical pest control options such as dormant oil would most likely provide little more than temporary, localized, control. The use of pesticides to control adelgid is not recommended in the riparian area due to potential impacts on water quality and aquatic life.

Without a major intervention (as yet unplanned), it is likely that the process of gradual infestation and demise of local hemlock stands by woolly adelgid will follow the patterns observed in areas already affected to the south. Reports from Southern Connecticut describe the re-colonization of hemlock sites by black birch, red maple and oak (Orwig, 2001). This transition from a dark, cool, sheltered coniferous stand to open hardwood cover is likely to raise soil temperatures and reduce soil moisture for sites where hemlocks currently dominate vegetative cover. Likewise, in the streams, water temperatures are likely to increase and the presence of thermal refuge for cool water loving fish such as trout, are likely to diminish. Alternatives for maintaining coniferous cover on hemlock sites include the planting of adelgid resistant conifers such as white pine as the hemlock dies out in the stand (Ward, 2001).

Human Disturbance and its Effects on the Riparian Vegetation

Although natural events disrupt growth and succession of riparian vegetation, human activities frequently transform the environment and, as a result, can have a long lasting impact on the capability of vegetation to survive and function. Presently, the most significant sources of human disturbance on riparian vegetation along the East Kill includes the construction and maintenance of roadway infrastructure, the maintenance of utility lines and the development of homes and gardens near the stream and its floodplain.

Roadway and Utility Line Influences on Riparian Vegetation

Due to the narrow valley and steepness of the valley walls, the alignments of County Routes 23C and 17 closely follow portions of the East Kill's stream alignment. Use and maintenance of the road right-of-way impacts the vigor of riparian vegetation. The narrow buffer of land between the creek and the road receives the runoff of salt, gravel, and chemicals from the road that stunt vegetation growth or increase its mortality. Road maintenance activities also regularly disturb the soil along the shoulder and on the road cut banks. This disturbance fosters the establishment of undesirable invasive plants. The linear gap in the canopy created by the roadway separates the riparian vegetation from the upland plant communities. This opening also allows light into the vegetative understory which may preclude the establishment of shade loving plants such as black cherry and hemlock.

Utility lines parallel the roadway and cross the stream at various points requiring the utility company to cut swaths through the riparian vegetation at each crossing, further fragmenting essential beltways for animal movement from streamside to upland areas. Although the road right-of-way and utility line sometimes overlap, at several locations along the stream, the right-of-way crosses through the riparian area separate from the road. This further reduces the vigor of riparian vegetation and prevents the vegetation from achieving the later stages of natural succession, typified by climax species such as sugar maple, beech and hemlock.



Utility crossing.

Residential Development Influence

Residential land use and development of new homes can have a great impact on the watershed and the ecology of the riparian area. Houses require access roads and utility lines that frequently have to cross the stream. Homeowners, who love the stream and want to be



Streamside development and limited riparian vegetation leads to compromised streambanks. This bank has been reinforced with riprap.

close to it, may clear trees and shrubs to provide access and views of the stream. Following this clearing, the stream bank begins to erode, the channel over-widens and shallows. The wide, shallow condition results in greater bedload deposition and increases stress on the unprotected bank. Eventually stream alignment may change and begin to cause erosion on the property of downstream landowners. Catskill stream banks require a mix of vegetation such as grasses and

herbs that have a shallower rooting depth, shrubs with a medium root depth, and trees with deep roots. Grasses alone are insufficient to maintain bank stability in steeply sloping streams such as the East Kill.

Many people live close to the stream and maintain access to the water without destabilizing the bank. By carefully selecting a route from the house to the water's edge and locating access points where the force of the water on the bank under high flow is lower, landowners can minimize disturbance to riparian vegetation and stream banks. Restricting access to foot traffic, minimizing disturbance in the flood prone area, and promoting a dense natural buffer provide property protection and a serene place that people and wildlife can enjoy. Additional information on concepts of streamside gardening and riparian buffers can be found at the following web site produced by the Connecticut River Joint Commission, Inc: <http://www.crjc.org/riparianbuffers.htm> and Catskillstreams.org. A list of native trees and shrubs "Native Trees for Riparian Buffers in the Upper Connecticut River Valley of New Hampshire and Vermont" developed by this group is provided in Appendix A. A list of native vegetation for the Catskill Mountain Region has been compiled using several sources, see Appendix A for this list or contact GCSWCD for more information.

Japanese Knotweed and Riparian Vegetation

Sometimes the attempt to beautify a home with new and different plants introduces a plant that spreads out of control and “invades” the native plant community. Invasive plants present a threat when they alter the ecology of the native plant community. This impact may extend to an alteration of the landscape should the invasive plant destabilize the geomorphology of the watershed (Malanson, 2002). The spread of Japanese knotweed (*Polygonum cuspidatum*), an exotic, invasive plant gaining a foothold in the East Kill, is an example of a plant capable of causing such a disruption. As its common name implies, Japanese knotweed’s origins are in Asia, and it was brought to this county as an ornamental garden plant.

Japanese knotweed is quite recognizable throughout the year. The series of photographs in Figure 2.7.3 illustrate different stages of Japanese knotweed’s growth throughout each season. This herbaceous, or non-woody, perennial goes through these cycles every year. In the spring (generally late April, early May), new red, asparagus-like shoots sprout from last year’s crown or from underground roots (*rhizomes*). By July, individual stems may reach 11 feet tall. Many thick, hollow stems are based at a crown.



Figure 2.7.3. Stages of Japanese knotweed’s growth throughout the growing season.

The upper areas of the stems form a few branches that reach out like an umbrella from the crown. Each main stem and branch holds several large, nearly-triangular leaves. In August knotweed dons abundant clusters of small, white flowers that attract several pollinators, such as bees, wasps and Japanese beetles.



Knotweed seeds.

The numerous flowers turn into buckwheat-like seeds by late September, early October. Although some seeds may create small seedlings (Forman & Kesseli 2003), knotweed spreads more by their *rhizomes*.

Cold weather halts the growth of knotweed; once frost covers the land, knotweed drops its leaves and turns an auburn hue. These dead stems often remain standing for one or two years and then cover the ground, decaying slowly.



Knotweed following a frost.



Knotweed's leaf pattern.



Dense stand of knotweed.

The above ground portion dies back each fall and re-grows to a height of 6-9 feet tall each spring. The canopy of the dense stands of bamboo-like stalks, covered by large heart shaped leaves, blocks out almost all light from reaching the soil, thereby shading out other plants and leaving the soil bare.

Japanese knotweed spreads primarily by vegetative means. Often, earthmoving contractors, highway department crews or gardeners transfer small portions of the roots in fill or soil that gets dumped on or near a streambank. These roots then grow into a new plant that soon becomes a colony. Japanese knotweed is able to spread rapidly on disturbed sites and prefers the moist, open conditions of the stream edge and bank for colonization. Once

knotweed has established itself in the riparian area, it is able to spread downstream after disturbances caused by beaver activity or by high flows scouring the streambank. Such disturbances often cause stems and rhizomes to break off and float downstream where fragments may establish themselves on streambanks that were previously unaffected by knotweed. Exposed streamside areas such as sediment deposits or disturbed banks with eroded soils lacking vegetation are particularly vulnerable to invasion by knotweed.

Although the impact of a Japanese knotweed invasion on the ecology of the riparian area is not fully understood, the traits of Japanese knotweed pose several concerns. Some of these concerns include:

- Knotweed appears to be less effective at stabilizing streambanks than deeper-rooted shrubs and trees, possibly resulting in more rapid bank erosion.
- The shade of its broad leaves and the cover by its dead litter limit the growth of native plants that provide food and shelter for associated native animals.
- Knotweed branches do not lean out over stream channels, providing little cooling from shade.
- Dead knotweed leaves (*detritus*) may alter food webs and impact the food supply for terrestrial and aquatic life.
- Large stands of knotweed impede access to waterways for fishing and streamside hiking.
- The presence of knotweed could reduce property value.
- Knotweed may alter the chemical make-up of the soil, altering soil microfauna and soil properties.

Japanese knotweed is very difficult to control. The broad use of herbicides, while potentially effective following a protocol of repeated treatments by a professional certified applicator, does present risks due to the threat the chemicals pose to water quality and the fragile aquatic ecosystem. Mechanical control, by



Japanese knotweed colony along the East Kill.

cutting or pulling, is labor intensive and requires regular attention to remove any re-growth. Biological controls are untested. The first step for residents and those who manage land and infrastructure in the East Kill is to familiarize themselves with the appearance and habits of knotweed. Next, it is important for landowners and land managers to monitor its spread. Landowners should avoid practices that would destabilize the stream banks or weaken the natural riparian vegetation that can prevent its spread. Any fill material introduced to the riparian area should be tested for the presence of Japanese knotweed. Any Japanese knotweed roots pulled or dug up from your property should be disposed of in a manner that will prevent it from spreading or re-establishing itself.

During the 2006 stream feature inventory and assessment, the project team mapped the distribution of Japanese knotweed along the East Kill. During these mapping efforts, the size of a colony was estimated; however the map does not show the area covered by each colony, only the presence of a colony (Figure 2.7.4). As is evident from the map, in 2006, Japanese knotweed was absent from more than half of




Figure 2.7.4. Japanese knotweed along the East Kill

the stream corridor; the first occurrence of knotweed was found in management unit #7, more than 8.5 miles downstream from the headwaters of the East Kill. Knotweed had colonized many sites along the East Kill downstream of its first occurrence, with 53 occurrences of knotweed affecting approximately 1,556 feet of stream banks (Figure 2.7.4). Without control efforts it is likely to continue to spread and fill in along the banks within a matter of a few years. Since much of the stream corridor has not been colonized by knotweed, it is critical to contain its spread and prevent invasion along unaffected stream banks. For more information about the specific quality and composition of a particular riparian area, please refer to individual Management Unit descriptions.

For several years, NYCDEP, GCSWCD and Hudsonia have been collaborating on research efforts to gain a greater understanding of Japanese knotweed. In 2003, Hudsonia submitted a final draft of their report *Japanese Knotweed and Water Quality on the Batavia Kill in Greene County, New York: Background Information and Literature Review*. This report provided a review of available information on Japanese knotweed including, the biology, ecology, history of its invasive spread, and methods of management (Talmage and Kiviat, 2003). Currently, the NYCDEP, GCSWCD and Hudsonia are working together on a project, along the Batavia Kill, to determine an effective treatment method for Japanese knotweed. Three experimental treatment methods are being tested for their effectiveness at managing stands of knotweed; these methods include, 1) frequent mowing, 2) limited excavation with planting of native species and a weed barrier around the trees, and 3) herbicide injection. This research is part of an effort to develop management recommendations for its future control. For more information on Japanese knotweed and for a link to Hudsonia's report, visit the GCSWCD website <http://www.gcswcd.com/stream/knotweed/> and/or Catskillstreams.org.

Japanese knotweed has established colonies along the East Kill and some of its' associated tributaries. Management of knotweed is a difficult task and careful consideration must be taken before determining how to manage individual knotweed stands or colonies on streamside properties. Besides understanding key characteristics about knotweed (e.g. how it spreads, what environments it prefers), it is also essential to recognize a few key concepts that actually apply to most invasive species. First and foremost, prevention is the best policy. Preventing the spread of knotweed is the most cost effective and time efficient approach to take, and may be achieved by: telling others about knotweed and warning them of its associated problems; keeping streambanks stable by allowing native trees and shrubs to grow and mature; and checking transported soil for any knotweed fragments.

It is critical to recognize that knotweed grows under diverse conditions and in varying locations, so there are different ways to approach its control. Before simply mowing down all the knotweed or spraying herbicides everywhere, one should first ask:

- How large is the stand of knotweed?
- Is it located near a waterway?
- What native plants exist nearby?

With answers to the questions above a customized approach may be taken, saving time and money by applying the most appropriate techniques.

Finally, someone wanting to control knotweed should understand that:

- **A disposal plan for all knotweed material is a must;** otherwise a new colony will just sprout somewhere else. This might include burning the material, burying it more than 6 ft. deep or letting it completely dry out.
- **Most treatments require multiple applications.** A one-time cutting or mowing of knotweed will not do anything besides stunt it temporarily and cause the rhizomes to extend underground faster towards more nutrients, possibly causing a higher rate of spread.
- **Revegetation with native species after treatment is necessary.** Leaving bare ground only promotes the reinvasion of knotweed. Rapid-growing, native trees and shrubs must be planted soon after removing knotweed in order to affect the most beneficial change.

Below are various treatment prescriptions depending on size of the knotweed stand, its proximity to a waterway and amount of surrounding vegetation. Please note that where bare ground exists after removing knotweed stems and roots, it is essential to revegetate the area with competitive (fast-growing) native trees and shrubs. This is especially critical if surrounding vegetation is limited or nonexistent. Otherwise reestablishment of knotweed is likely and control efforts are futile.

For *small* stands (less than 3ft²):

- Cover with dark plastic.
- Frequent cutting, grubbing or pulling with safe disposal of knotweed stems.
- Herbicide injection of stems. PLEASE READ HERBICIDE CAUTION BELOW.

For *medium* stands (3ft² to 25ft²):

- Frequent mowing (do not allow cut material to leave site).

For *large* stands (25ft²+):

In some cases, the extent of a knotweed colony is so extensive that more harm (e.g. damage to soils) would be done in trying to eliminate the entire stand. For this reason control of expansion is the appropriate action.

- Frequent mowing around edges of stand (do not allow cut material to leave site).
- Herbicide injection of stems in edges of stand. PLEASE READ HERBICIDE CAUTION BELOW.

Herbicide Caution: Glyphosate (e.g. Rodeo, Roundup, Aquamaster) is the recommended active agent. When used with care and according to product labels, this herbicide does NOT negatively affect *untouched* plants and animals. Using an injection method is the highest recommendation, because knotweed material is not cut therefore requiring no disposal. Also this method eliminates drift and targets only injected stems. Only certain herbicides, such as Rodeo and Aquamaster, should be used near a waterway. Please take care to wear appropriate protective equipment. Check with Cornell Cooperative Extension of Greene County at 518-622-9820 for information about the proper, safe and legal use of herbicides.

Assessment of the Current Condition of Riparian Vegetation

As part of the stream management planning process, physiognomic classes (e.g., deciduous open tree canopy, shrub land, herbaceous) were mapped and the riparian vegetation assessed for the East Kill watershed (methodology available in Appendix B). The purpose of this exercise was to provide the planning team with baseline information about communities present in the watershed, a description of the condition of vegetation in the riparian area, and to aid in the development of recommendations related to the management of riparian vegetation along the stream.

Mapping of Physiognomic Classes

Mapping of physiognomic classes was loosely based on the Vegetation Classification Standard produced by The Federal Geographic Data Committee. The mapping was based upon 2001 digital-ortho photos and confined to the riparian and near adjoining upland areas within 300 ft. of the mainstem of the East Kill. This classification was selected because it allows identification of locations, such as herbaceous or cobble deposits, where the



Riparian vegetation (closed mixed), protects stream banks and water quality, and provides habitat and food sources for fish and wildlife.

combination of channel morphology and riparian vegetation would indicate the greatest cost-benefit from riparian buffer plantings and bio-engineered bank stabilizations.

The mapping exercise included the approximate delineation of the classes through the photointerpretation of 2001 infra-red digital orthophotography acquired by New York State. A physiognomic class GIS data layer was created using heads-up digitizing techniques with ESRI's Arcview software. The photo interpretation was field checked with class boundaries, and classifications were amended based upon field observations. The vegetation map resulting from this process is folded and included in the back of this management plan.

Summary of Findings

According to this riparian vegetation assessment, evergreen closed tree canopy (approximately 289 acres) and herbaceous (approximately 231 acres) were the largest physiognomic classes within the 300ft. buffer, while mixed closed tree canopy and deciduous closed tree canopy occupied approximately 206 acres and 167 acres respectively. This predominance of closed forest cover helps to provide a high degree of stability to the watershed. Forest cover slows storm runoff, and the stream banks in much of the watershed have some woody vegetation to protect against bank erosion. Protection of the forest communities near the stream will help ensure long-term stream stability, but the effectiveness of stream protection provided by vegetative communities, differs based on their width, plant density, vegetation type and the stream's geomorphic characteristics.

Although forested land covered a large portion of the watershed's riparian area, the extent of herbaceous cover is a concern. While herbaceous cover is better than no cover at all, plants with a variety of rooting depths (herbs, shrubs and trees) provide more extensive stream bank protection. Approximately 262 acres, or 21% of land area was considered to have inadequate vegetative cover; this included areas of herbaceous vegetation, bare soil and revetment. A streamside planting program is recommended to address these areas of inadequate vegetation.

Table 2.7.1 provides the results of the GIS vegetation assessment of the East Kill, including the area and percentage of each land cover type. Classes listed in italics contribute to the total area of inadequate vegetation.

Table 2.7.1. Summary of Physiognomic Vegetation Classification		
Vegetation Classification	Area (acres)	Percent
<i>Bare Soil</i>	28.10	2.22%
Deciduous Closed Tree Canopy	166.55	13.19%
Deciduous Open Tree Canopy	85.78	6.79%
Evergreen Closed Tree Canopy	288.96	22.88%
Evergreen Open Tree Canopy	19.90	1.58%
<i>Herbaceous Vegetation</i>	231.48	18.33%
Impervious Surface	28.63	2.27%
Mixed Closed Tree Canopy	205.57	16.28%
Mixed Open Tree Canopy	24.76	1.96%
<i>Revetment</i>	2.52	0.20%
Shrubland	93.22	7.38%
Unpaved Road	5.89	0.47%
Water	81.51	6.45%
Total Area	1262.87	
Inadequate Vegetation	262.09	20.75%

Riparian ecosystems are an important component of watershed protection and resource conservation. Therefore, it is important to maintain and improve the riparian vegetation along the East Kill and its' tributaries.

East Kill Streamside Planting

A streamside planting program is recommended for the East Kill. The findings from the mapping of physiognomic classes can be used to identify candidate stream reaches for inclusion in future streamside planting programs. There are three main steps to establish this type of program.

1. *Identify priority sites* using information gathered during riparian vegetation characterization analyses to identify potential planting sites where improvement of the riparian vegetation is likely to be both effective and successful.
2. *Develop treatment designs* for participating prioritized sites using primarily native plants that address landowner aesthetics, ecological enhancement and water quality improvement or protection.
3. *Install the designs* and document the planting process and results for program replication and general education/outreach.

To effectively carry out riparian planting projects, it is necessary to develop objective physical criteria for identifying and prioritizing eroding banks that may be stabilized with riparian vegetation plantings. It is also important to coordinate with streamside landowners by canvassing riparian landowners whose properties meet these objective criteria for their interest in participating in a project, and establishing a partnership with interested landowners. Technical assistance may then be provided to landowners to reestablish the riparian buffer on their property with native vegetation. GCSWCD and NYCDEP will be piloting such a project in 2007.

References:

- Adams, M.S., Terzilla, D. & Baum, B.S. (2002) Community-based monitoring in the Catskills. In B. Onken, R. Reardon, J. Lashomb (Eds.), *Symposium on the hemlock woolly adelgid in eastern North America: February 5-7, 2002, East Brunswick, New Jersey* (pp. 100-105). New Brunswick, NJ: USDA Forest Service – Rutgers University.
- Forman, J. & Kesseli, R.V. (2003). Sexual reproduction in the invasive species *Fallopia japonica* (Polygonaceae)¹. *American Journal of Botany* 2003, 90, 586-592.
- Kudish, M. (2000). *The Catskill forest: a history*. Fleischmanns, NY: Purple Mountain Press, Ltd.
- Malanson, G.P. (1993). *Riparian landscapes*. Cambridge, UK: Cambridge University Press.
- McClure, Mark S. and Carole A., and Cheah, S-J., 2002, Important mortality factors in the life cycle of hemlock woolly adelgid, *Adelges tsugae* Annand (Homoptera: Adelgidae). In B. Onken, R. Reardon, J. Lashomb (Eds.), *Symposium on the hemlock woolly adelgid in eastern North America: February 5-7, 2002, East Brunswick, New Jersey* (pp. 13-22). New Brunswick, NJ: USDA Forest Service – Rutgers University.
- Orwig, D.A. (2002). Stand dynamics associated with chronic hemlock woolly adelgid infestations in southern New England. In B. Onken, R. Reardon, J. Lashomb (Eds.), *Symposium on the hemlock woolly adelgid in eastern North America: February 5-7, 2002, East Brunswick, New Jersey* (pp. 36-46). New Brunswick, NJ: USDA Forest Service – Rutgers University.
- Talmage, E. & Kiviat, E. 2003. *Japanese knotweed and water quality on the Batavia Kill in Greene County, New York: background information and literature review*. Annandale, NY: Hudsonia Ltd.
- Ward, J. (2002). Restoration of damaged stands: dealing with the after effects of hemlock woolly adelgid. In B. Onken, R. Reardon, J. Lashomb (Eds.), *Symposium on the hemlock woolly adelgid in eastern North America: February 5-7, 2002, East Brunswick, New Jersey* (pp. 118-126). New Brunswick, NJ: USDA Forest Service – Rutgers University.

2.8 Land Cover/Land Use

Land use and land cover of a watershed have a great influence on water quality and stream stability. The watershed's land cover directly impacts stream hydrology by influencing the amount of stormwater runoff. Forests, natural meadows and wetlands naturally absorb rainwater, allowing a portion of it to percolate back into the ground. However, impervious surfaces such as pavement, parking lots, driveways, hard-packed dirt roads and rooftops increase the amount of rainfall that flows over land and reduces the amount of rainfall that percolates into the soil to recharge groundwater wells and streams.

Impervious cover is a major influence on streams and stream life due to the way it changes the amount and duration of stormwater that gets to the stream. Generally, the more impervious surface there is in a watershed, the less groundwater recharge (which supplies summer low flows), and the greater the magnitude of storm flows (and related erosion in streambeds). In addition to degrading streams, watersheds with a high percentage of impervious surfaces are prone to larger and more frequent floods, which cause property damage through inundation, as well as ecological harm resulting from lower base stream flows.

The literature has documented the deleterious effects impervious surfaces have on biota (Limburg and Schmidt, 1990; May et al., 2000; Wang et al., 2001; Roy et al., 2005), stream stability (Booth, 1990; CWP, 1998; White and Greer, 2005; Wohl, 2005) and in-stream water quality (Groffman et al., 2004 and Deacon et al., 2005). For example, impervious surfaces can raise the temperature of stormwater runoff, which in turn reduces the water's ability to hold dissolved oxygen and harms some game fish populations, while promoting excess algal growth. Field observation, research and hydrologic modeling suggest a threshold of 10% impervious surface in a watershed, after which there is marked transition to degraded stream conditions (CWP, 1998 and Booth, 2000).

Certain types of pollution problems are often associated with particular land uses, such as sedimentation from construction activities. There has been a vast array of research demonstrating that as land uses become more urbanized (built), biotic communities decline in health (Limburg and Schmidt, 1990; Schueler and Holland, 2000; May et al., 2000; Wang et al., 2001 and Potter et al. 2005). Concentrations of selected chemical constituents, including

nitrate, in stream base-flow were strongly affected by the predominant land use in a large Hudson Valley study (Heisig, 2000). The decline of watershed forest cover below 65% percent marked a transition to degraded water quality (Booth, 2000). Based upon these results, land use/cover appear to be attractive attributes for long-term trend tracking. These results can then be correlated with in-stream water quality data and then used to focus best management practices towards the land uses with the greatest impact on water quality.

Land cover of the East Kill Watershed was analyzed using the LANDSAT ETM 2001 geographic information system (GIS) coverage created by the New York City Department of Environmental Protection (DEP) (Figure 2.8.1). The categories are comprised of 47 different classification descriptions. To simplify the data, land cover classifications have been grouped together and re-classified to convey the general land cover category that each classification falls



Figure 2.8.1. East Kill watershed

under. For example, the classification descriptions of central business district, residential, and industrial, among others, have been combined and re-classified as development.

In 2001, approximately 90% of the East Kill watershed was covered by coniferous, deciduous, or mixed forest, while development covered approximately 2.5% (Table 2.8.1 and Figure 2.8.2). Proper land use planning to direct development into priority areas while preserving sensitive areas should be utilized to limit the impact of future development and subsequent increases in impervious surfaces.

Land Cover Category	Acres	Percent Cover
Agriculture	228.26	0.98%
Development	582.04	2.51%
Dumps	0.01	0%
Exposed Soil	0	0%
Forest	20,808.70	89.71%
Herbaceous	191.34	0.83%
Managed Herbaceous	247.89	1.07%
Mined Lands	0	0%
Shrubland	684.25	2.95%
Water	103.97	0.45%
Wetland	348.60	1.50%
Total	23,195	100%




Figure 2.8.2. Land Cover of the East Kill Watershed in 2001. Large format map is available in the back pocket of this plan.

Protected Lands

To determine the percentage of parcels within the East Kill basin that were protected as Wild, Forested, Conservation Lands & Public Parks, the ownership and property use classifications as documented on records of the Greene, Schoharie and Delaware County Real Property Tax Service Departments, were analyzed. In 2006, approximately 34% of the East Kill Watershed lands were protected as Wild, Forested, and Conservation Lands & Public Parks. The primary owner of the protected lands was New York State with ownership of 82% of protected land in the East Kill Watershed. Under current State laws, these lands owned by the State will remain undeveloped. In 2006, approximately 0.3% of land within the Watershed was owned by New York City (Table 2.8.2).

Property Use Class		
	Acres	Percent
Wild, Forested, Conservation Lands & Public Parks	7,714	33.8%
NYC Owned Land	64	0.3%

References:

- Booth, D.B. 1990. Stream Channel Incision Following Drainage Basin Urbanization. Water Resources Bulletin Volume 26: 407-417.
- Booth, D. 2000. Forest Cover, Impervious Surface Area, and the Mitigation of Urbanization Impacts in King County, Washington. Center for Urban Water Resources Management, University of Washington, Seattle, WA.
- CWP. 1998. Rapid Watershed Planning Handbook: A Comprehensive Guide for Managing Urbanizing Watersheds. Center for Watershed Protection, Ellicott City, Maryland.
- Deacon, J.R., Soule, S.A., and Smith, T.E. 2005. Effects of Urbanization on Stream Quality at Selected Sites in the Seacoast Region in New Hampshire, 2001-03. United States Geological Survey Investigations Report 2005-5103, 18 p.
- Groffman, P.M., Law, N.L., Belt, K.T., Band, L.E., and Fisher, G.T. 2004. Nitrogen Fluxes and Retention in Urban Watershed Ecosystems. Ecosystems 7: 393– 403.
- US EPA, 2003 - Land Use Planning: <http://www.epa.gov/watertrain/protection/r2.html>

- Heisig, P. 2000. Effects of Residential and Agricultural Land Uses on the Chemical Quality of Baseflow of Small Streams in the Croton Watershed, Southeastern New York. Publication # WRIR 99-4173. United States Geological Survey, Troy, NY.
- Limburg, K.E. and Schmidt, R.E. 1990. Patterns of Fish Spawning in Hudson River Tributaries: Response to an Urban Gradient?. Ecology Volume 71 (4): 1238 – 1245. 05).
- May, C.W., Horner, R.R., Karr, J.R., Mar, B.W. and Welch, E.B. 2000. Effects of Urbanization on Small Streams in the Puget Sound Ecoregion. Watershed Protection Techniques, 2(4): 483-494.
- Potter, P.M., Cabbage, F.W., and Schaberg, R.H. 2005. Multiple-scale landscape predictors of benthic macroinvertebrate community structure in North Carolina. Landscape and Urban Planning 71: 77-90.
- Roy, A. H., Freeman, M. C., Freeman, B. J., Wenger, S. J., Ensign, W. E., Meyer, J. L. 2005. Investigating hydrologic alteration as a mechanism of fish assemblage shifts in urbanizing streams. Journal of the North American Benthological Society Volume 24 (3): 656-678.
- Scheuler, T.R. and Holland, H.K. 2000. Housing Density and Urban Land Use as Indicators of Stream Quality. In: The Practice of Watershed Protection 2(4): 735-739.
- Wang, L., Lyons, J., Kanehl, P. and Bannerman, R. 2001. Impacts of Urbanization on Stream Habitat and Fish Across Multiple Spatial Scales. Environmental Management Vol. 28(2): 255-266.
- White, M.D. and Greer, K.A. 2005. The effects of watershed urbanization on the stream hydrology and riparian vegetation of Los Peñasquitos Creek, California. Landscape and Urban Planning, Volume 74(2): 125-138.
- Wohl, E. 2005. Compromised Rivers: Understanding Historical Human Impacts on Rivers in the Context of Restoration. Ecology and Society 10(2): Article 2.

2.9 Wildlife and Fisheries

The East Kill watershed is literally crawling with life. An amazing variety of habitats, people, plants, and animals are all interconnected in a fragile web of life, often called biodiversity. Every member is essential to keeping this web in balance. For example, the list of species required for the life cycle of a single tree may be in the hundreds or thousands. Moreover, the list of animals that will utilize a single fallen tree is



A very young fawn crosses the Schoharie Creek, summer 2006.

in the thousands, but a few of the more well known creatures include squirrels, woodpeckers, grouse, bears, foxes, skunks, mice, and shrews as well as worms, salamanders, beetles, ants, centipedes, sowbugs, and other insect larvae. There are twice as many species of beetles that live on dead and dying wood as there are species of mammals, birds, reptiles, and amphibians in the entire world (Kyker-Snowman, 2003). The fallen tree also provides critical habitat, steady moisture, and food for a multitude of mosses, fungi, trees, and vascular plants. If our fallen tree had been removed either during land use changes or during “clean up” efforts after falling, the ramifications would reverberate throughout the web. Certainly, this doesn’t preclude us from taking a few trees for firewood, but if enough fallen trees are removed, the structure of the overall community would likely change.

The fallen tree example was meant to demonstrate the complexity of the web of life, and how eliminating one organism or habitat will ultimately affect many. It is very difficult to predict the consequences of removing individual strands from the web of life. Therefore, as an integral piece of the web, humans should work toward protection and preservation of the functions necessary for our survival. There are many ecosystem functions we receive from nature including cleaner air through vegetation respiration, cleaner water through soil and wetland filtration, soil formation from forests, pollination of food crops from our native insects, natural flood water retention/groundwater recharge, and pest control from our native bats, birds, and insects (e.g. dragonflies/damselflies). For example, bees pollinate about a

trillion apple blossoms each year in New York State, micro-organisms biodegrade much of our garbage as well as fallen leaves, sticks and other dead animal and plant matter, soil bacteria turn nitrogen into nitrate fertilizer and plants use up carbon dioxide and produce oxygen, thereby slowing global climate change. One example that affects us locally is the maintenance of healthy biodiversity and community structure, which if done properly can reduce the incidence of Lyme disease (LoGiudice et al., 2003); and forest fragmentation which can increase white-footed mouse populations, that in turn increases the human risk of exposure to Lyme disease (Allan et al. 2003). Therefore, the benefits of a healthy and diverse ecosystem extend far beyond clean air and water and into the fabric of human health and quality of life.

In the United States the economic services provided by a vibrant/healthy biological web of life (biodiversity) contribute an estimated \$319 billion per year, or 5% of the gross domestic product (Pimentel et al., 1997). The worldwide benefits are estimated to be \$2,928 billion per year, or approximately 11% of the world economy (Pimentel et al., 1997). Closer to home, the economic impact of the East Kill hasn't been calculated, but anecdotally considering it's uses for recreation, water supply and aesthetics the economic value is most likely high. Clearly, our economic vitality depends on maintaining healthy biodiversity, which in turn indicates clean water and a good quality of life.

The plants and animals that inhabit the East Kill watershed are suited to the habitats provided by our temperate climate. The other major factor is human alteration of the landscape. Pre-European colonization the watershed was predominantly forested with some small areas cleared by Native Americans for hunting. Early European settlers attempted to farm the land, but abandoned it soon after due to a short growing season, steep slopes and rocky and shallow



Milk snake in the East Kill, summer 2006.

soils. Between 1800 and the early 1900s gristmills, woolen mills, sawmills, the tanning industry, quarrying for bluestone, logging, furniture making, railroads and resorts cleared the

Catskills of its forest cover. Since the early 1900s these industries have declined and areas that were previously cleared have grown back into forest, with approximately 90% of the East Kill basin's land cover being classified as forest in a 2001 NYCDEP analysis. The reaction of wildlife has varied to the changing land uses. A few, such as the timber wolf, eastern cougar, New England cottontail and passenger pigeon have been extirpated from the region (passenger pigeon is extinct worldwide); and some such as tiger beetle and timber rattlesnake are disappearing from the Catskills. Beaver, pileated woodpeckers, and bald eagles were once gone from this region due to over hunting, habitat loss, and pesticide poisoning respectively, but have since returned with reduced hunting pressure, an increase in second-growth forests, and a ban on DDT. Some species, such as the bobcat, black bear and osprey are less common than they were prior to European colonization. However, other common species, such as the white tailed deer, raccoon, skunk, red fox, robin, and painted turtle have thrived.



Green frog in the East Kill, summer 2006.

We often focus on human-induced land use changes as the dominant factor in habitat and natural landscape changes. However, many wildlife and plant species also influence the landscape. Heavy deer browsing of seeds, seedlings, and saplings can dramatically alter the composition of a forest to encourage the growth of species that deer find less palatable (Curtis, 2004). Species imported from other areas that thrive in our region, often called invasive species, can also have dramatic effects on the landscape. For example, Japanese knotweed (*Polygonum cuspidatum*) is native to Asia, but has run rampant in the Schoharie basin choking out native species, diminishing recreation opportunities and possibly making soil more susceptible to erosion (more info in section 2.6).



Tent caterpillars along the Schoharie Creek, summer 2006.

The woolly adelgid (*Adelges tsugae*), a small aphid-like insect pest native to China and Japan, is threatening to decimate our eastern hemlock (*Tsuga canadensis*) populations. Once infested, hemlock mortality rates range between 50%-99% (Orwig, 2002). The plant species most likely to replace hemlocks are hardwood tree species and possibly other invasive species. Ultimately, this will have a dramatic effect on the structure of these communities. For example, the distribution and abundance of brook trout and diversity of aquatic insects will likely decline with the hemlock forests (Evans, 2002). Hemlock forests maintain stable, lower water temperatures and more stable hydrologic regimes (i.e. they don't dry up as much) than the hardwood forests that will likely replace them (Snyder et al., 2002). These are just a few examples of how, in a global society, careless actions can import and release species that can drastically change our ecological communities.

Native pests often have native predators that control their populations. For example, the forest tent caterpillar (*Malacosoma disstria*) can cause a large amount of damage to Catskill forests. However, their population tends to be controlled by a natural predator fly (*Sarcophaga aldrichi*) whose populations explode following the explosion of the caterpillar's population and help bring the caterpillar populations back under control. A bacterial disease, known as "wilt" and cold, wet, weather conditions in early spring also help to control the caterpillar population. This demonstrates the checks-and-balances of native versus non-native pests. Native pests often have a naturally-evolved control measure that eventually bring the populations under control, but non-native species do not.

The East Kill is primarily a cold water stream, meaning it provides suitable water temperatures for organisms, such as brook trout and sculpins, which require cold water (less than 72° F (22°C)). The East Kill is stocked annually with 2,700 brown trout from its confluence with the Schoharie upstream approximately 10.3 miles (NYSDEC, 1993). Upstream of this point (~2.4 miles downstream



Tent caterpillar damage within the Schoharie Watershed, summer, 2006. Favorable climatic conditions were good, so many of these trees probably grew a second growth of leaves after the caterpillar population dwindled in late June.

of Colgate Lake) to its source the East Kill is considered to have a sufficient wild trout population to make stocking unnecessary. The wild trout population is comprised primarily of brown trout to Harriman Lake, and brook trout upstream of Harriman Lake. Fish communities were analyzed approximately halfway through the East Kill from headwaters to mouth (near the Route 40/23C intersection) in 2000, 2002 and 2003. A lumped summary (3 years worth of data) of all the fish species collected is available in Table 2.9.1. Interestingly, researchers found that over a relatively short period of time (3 years), modified-natural channel design restorations that incorporated fish habitats increased fish biomass, including trout biomass and numbers (Baldigo et al., 2006).

Table 2.9.1. Fish species present in the East Kill in 2000, 2002 and/or 2003 and their relative abundance to each other (Personal communication, Barry Baldigo, 2007).

Common Name	Scientific Name
Creek chub	<i>Semotilus atromaculatus</i>
Common shiner	<i>Luxilus cornutus</i>
Pumpkinseed	<i>Lepomis gibbosus</i>
Bluegill	<i>Lepomis macrochirus</i>
White Sucker	<i>Catostomus commersoni</i>
Stone Roller	<i>Campostoma anomalum</i>
Cutlips minnow	<i>Exoglossum maxillingua</i>
Marginated madtom	<i>Noturus insignis</i>
Brown trout	<i>Salmo trutta</i>
Brook trout	<i>Salvelinus fontinalis</i>
Blacknose dace	<i>Rhinichthys atratulus</i>
Longnose dace	<i>Rhinichthys cataractae</i>
Tesselated Darter	<i>Etheostoma olmstedii</i>
Largemouth Bass	<i>Micropterus salmoides</i>
Slimy Sculpin	<i>Cottus cognatus</i>
Stone Cat	<i>Noturus flavus</i>

Despite being a cold water fishery, it should be noted that a high-predatory warm-water species was cited often during the 2006 assessment. These largemouth bass were stocked or escaped into the stream from adjacent ponds and could compete with trout for resources. However, they may also migrate to more suitable habitat conditions and/or suffer high mortality rates during the cold winter months.



Largemouth bass in the East Kill, summer 2006.

Wildlife of Stream Corridors and Conservation Recommendations for the Upper Schoharie Watershed, Greene County, March, 2007

The Upper Schoharie Watershed contains a high degree of biological diversity with a species assemblage that is unusual within the Hudson River Valley. Forests with features such as talus slopes, cliffs, and mature stands are habitat for plants and animals adapted to these conditions. The large, unfragmented nature of the forests creates favorable habitat for wide-ranging animals (such as black bear and bobcat) and wildlife that prefer forest interiors (such as black-throated blue warbler). It is likely that forests of the Upper Schoharie watershed are important breeding areas for raptors such as broad-winged hawk, Northern goshawk, and sharp-shinned hawk.

Forests that occur adjacent to the stream create habitat for a wide range of small mammals, including rarely seen moles, voles, and shrews, and fox, weasel, mink, beaver, and muskrat. The change in elevation from stream valley floor to Catskill peaks, and the presence of both evergreen and deciduous forests contribute to the watershed's biodiversity. High-altitude coniferous forests are habitat for the rare Bicknell's thrush and blackpoll warbler.

In the Upper Schoharie watershed, abundant streams with cobble beds, undercut banks, and streamside wetlands and forests are habitat for damselflies, dragonflies, stream salamanders, turtles, and frogs. The wood turtle lives almost exclusively in and near streams, while spotted turtles might be found in streamside wetlands. Riparian forests are particularly important breeding habitat for birds such as the Louisiana waterthrush and yellow-throated

vireo. Stream corridors are the preferred foraging habitat for the many bat species that are likely to occur in the watershed.

Grassy fields, open woods, and shrubby patches make important contributions to biodiversity of the watershed. These open and scrubby areas can provide nesting habitat for the wood turtle and shrubland bird species that are declining in New York State as old farms revert to forests. Young forests are habitat for Canada warbler, American woodcock, and ruffed grouse, while open shrublands and dense thickets are preferred by brown thrasher.

Many species, like American woodcock and wood turtle, require a complex of different habitats to complete breeding, foraging, overwintering, and migration portions of their life cycles. As a result, maintaining connectivity between the stream and the adjacent uplands is very important for biodiversity conservation. NYSDEC Species of Greatest Conservation Need (SGCN), included in the State Wildlife Plan, and Hudson Valley regionally rare species that may use the Schoharie basin are listed in Table 2.9.2. A complete list of potential species and an occurrence map is provided in Appendix C, and a map of rare species and significant ecosystems is provided in Figure 2.9.1.

Management Recommendations

Stream managers should consider the following general recommendations to maintain and protect important stream corridor habitats:

- Limit disturbance and protect both small and large stream corridor wetlands that provide significant habitat for amphibians, reptiles, and breeding birds in the watershed;
- Most shrubland breeding birds are relatively tolerant of human development if appropriate habitats exist, and unlike some grassland birds, do not require large habitat patches for breeding. While open lands should not be created at the expense of mature, unfragmented forests, agricultural and suburban landowners who maintain shrubby thickets in the uplands adjacent to stream corridors can support shrubland birds;
- Where possible, plant native species appropriate to the pre-existing or predicted ecological community for a site;

- Stream managers are encouraged to learn to recognize the Appalachian tiger beetle and other declining and threatened species and report observations to the NY Natural Heritage Program.

Riparian buffer widths can be established to conserve habitat function, in addition to water quality, hydrologic, and geomorphic functions. It is particularly important to maintain habitat connectivity needed by wildlife to complete their life cycles. To evaluate connectivity, consider the needs of indicator species, or species of conservation concern in the watershed.

- The forest area within 300 ft of the forest edge is considered “edge” habitat. Edge habitats support increased densities of deer and invasive plants, and are avenues for nest predators to enter forests. A minimum 300 ft forested stream buffer will protect forest health and provide better breeding habitat for forest wildlife;
- Riparian forests at least 50 acres in size with an average total width of at least 300 ft can provide forest interior habitat and should be highly valued. Breeding bird diversity increases substantially between 300 and 1,500 ft from the stream’s edge;
- Most of the amphibian and reptile observations in this watershed are within or near stream corridors. Seek to create a minimum 500 ft forested buffer around stream corridor wetlands to provide terrestrial habitat required by stream- and vernal pool-breeding amphibians to complete their life cycles, and to protect wetlands from adjacent land uses;
- Stream buffers up to 1,000 ft will encompass most wood turtle nesting sites and summer habitats (wood turtles are on land during the spring and summer and over-winter in rivers). These buffers should be maintained along one or more miles of stream length to accommodate the yearly movements of wood turtles up and down the stream channel;
- Buffer widths of 30-100 ft should be maintained for riparian forest canopies to provide enough shading and cooling of streams to maintain trout populations. These buffers need to be nearly continuous. Some studies suggest 80% of banks

along a stream supporting trout populations must have forests at least 30 ft wide to provide sufficient shade for trout;

- Minimum buffers of 50-100 ft are often recommended to protect aquatic communities. Large woody debris deposited into streams provides important shelter for fish, and in particular for trout. At a minimum, a 50 ft buffer appears necessary to maintain sufficient woody debris inputs to streams. Riparian vegetation provides leaves and other forms of litter that feed macroinvertebrates. In turn, aquatic macroinvertebrates are the major food source for most freshwater fish. A minimum 100 ft buffer is recommended to protect aquatic macroinvertebrate and fish abundance.

Typically, the locations of wood turtle nesting sites are not known. However, stream managers can use the following information to identify possible nesting areas near the stream and maintain adequate buffer widths to protect nesting wood turtles:

- Wood turtles typically nest in sandy, bare areas well exposed to sunlight and close to water, but elevated. Usually, nesting areas are within 200 ft of the stream channel, but wood turtles will travel up to 2,000 ft from the stream to reach suitable nesting areas. Nesting sites are often exposed stream banks, but can include cultural features such as nearby railroad tracks, abandoned sand and gravel mines, utility right-of-ways, and meadows/fields with gravelly soils. Wood turtles will nest in corn fields and other recently disturbed areas. If possible, identify potential wood turtle nesting sites near streams and protect them. Buffers between the stream and suitable nesting areas should be used lightly or not at all, particularly during nesting season (May to July, peak in June);
- Where the wood turtle is likely to occur, maintain stream geomorphology with naturally elevated banks and gravel deposits (used for winter hibernation, basking, and nesting).

A number of stream corridor species depend on natural channel processes to provide habitat during parts of their life cycles:

- Stream salamanders are generally sensitive to siltation, scouring, nutrient enrichment, channelization, and diversion of water. Maintain natural stream processes and riparian buffers to protect salamander habitats. Spring salamander is probably the most sensitive species and is found in relatively unpolluted headwater streams of the Catskills;

- There is one known remaining occurrence of Appalachian tiger beetle in this watershed. There are only 10 rivers in NYS with populations of this species. The Appalachian tiger beetle (right) is



typically found on riverside sand and cobble bars at the edges of forested streams. Stream management practices should maintain natural stream processes, including natural flooding regimes that prevent dense plant growth on cobble bars. Gravel mining and off-road vehicle use of sand and gravel bars can destroy beetle larvae.

Table 2.9.2. NYSDEC Species of Greatest Conservation Need and other species of conservation concern that may occur in the Upper Schoharie watershed (a complete list of species and NYNHP classification descriptions are available in appendix C) (Prepared by the NYSDEC Hudson River Estuary Program, Feb. 2007).

Predicted Terrestrial Vertebrate Species (source: Hudson River Valley GAP)

Terrestrial, vertebrate species that are predicted to occur within the watershed based upon presumed associations of species with habitats. See the HRV-Gap Analysis Project report to view predicted species distribution maps.

Mammals		TNC Status	Legal Status	NYNHP Species	NYSDEC SGCN	Regional
Eastern Pipistrelle	Pipistrellus subflavus	G5, S3				R
Eastern Red Bat	Lasiurus borealis	G5, S5B, SZN			X	R
Hoary Bat	Lasiurus cinereus	G5, S4B, SZN			X	R
Indiana Myotis	Myotis sodalis	G2, S1	FE, SE	X	X	
Silver-haired Bat	Lasionycteris noctivagans	G5, S4B, SZN	SC		X	Rm
Woodland Jumping Mouse	Napaeozapus insignis	G5, S5				R
Long-tailed Shrew	Sorex dispar	G4, S4				R
Southern Bog Lemming	Synaptomys cooperi	G5, S4				R
Porcupine	Erethizon dorsatum	G5, S5				R
Black Bear	Ursus americanus	G5, S5	G			S

Fisher	<i>Martes pennanti</i>	G4G5, S4	G			S
Bobcat	<i>Lynx rufus</i>	G5, S4	G			V
Eastern Cougar	<i>Felis concolor cougar</i>	G5TH, SX	FE, SE		X	
River Otter	<i>Lutra canadensis</i>	G5, S5	G		X	S
Amphibians		TNC Status	Legal Status	NYNHP Species	NYSDEC SGCN	Regional
Jefferson Salamander	<i>Ambystoma jeffersonianum</i>	G5, S4	SC		X	
Jefferson Salamander Complex	<i>Ambystoma jeffersonianum x laterale</i>	G4, S3	SC		X	
Spotted Salamander	<i>Ambystoma maculatum</i>	G5, S5				V
Northern Dusky Salamander	<i>Desmognathus fuscus</i>	G5, S5				D, V
Longtail Salamander	<i>Eurycea longicauda longicauda</i>	G5, S2S3	SC	X	X	
Four-toed Salamander	<i>Hemidactylium scutatum</i>	G5			X	
Common Mudpuppy	<i>Necturus maculosus</i>	G5, S4			X	
Northern Red Salamander	<i>Pseudotriton ruber ruber</i>	G5, S3S4			X	
Northern Leopard Frog	<i>Rana pipiens</i>	G5, S5	G			R
Wood Frog	<i>Rana sylvatica</i>	G5, S5	G			V
Reptiles		TNC Status	Legal Status	NYNHP Species	NYSDEC SGCN	Regional
Wood Turtle	<i>Clemmys insculpta</i>	G4, S3	SC, G		X	
Timber Rattlesnake	<i>Crotalus horridus</i>	G5, S3	ST		X	
Smooth Greensnake	<i>Opheodrys vernalis</i>	G5, S4			X	D
Eastern Box Turtle	<i>Terrapene c. carolina</i>	G5, S3	SC		X	

Observed Breeding Birds (source: 1980-1985 Breeding Bird Atlas)

Breeding bird species known or suspected to be breeding within the watershed. The species list is derived from reports of observed breeding bird activity within Breeding Bird Atlas Blocks that overlap the watershed. Parties using these data for environmental review purposes do so at their own risk.

Key: TNC Status: For State and Global Rank explanations see www.natureserve.com; Legal Status: FE = Federal Endangered; FT = Federal Threatened; SE = State Endangered; ST = State Threatened; G = Game species; SC = State Special Concern; NYNHP Species: Rare species tracked by the NY Natural Heritage Program; NYSDEC SGCN: Species of Greatest Conservation Need included in State Wildlife Plan; Other Ranking: Listed Partners in Flight WatchList

Birds		TNC Status	Legal Status	NYNHP Species	NYSDEC SGCN	Other Ranking	# Blocks (35 total)
American Black Duck	<i>Anas rubripes</i>	S4, G4	G		X	WL, PIF	3
American Woodcock	<i>Scolopax minor</i>	S5, G5	G		X	WL, PIF	9
Bicknell's Thrush	<i>Catharus bicknelli</i>	S2S3B, G4	SC	X	X	WL, PIF	5
Black-billed Cuckoo	<i>Coccyzus erythrophthalmus</i>	S5, G5			X		2
Black-throated Blue Warbler	<i>Dendroica caerulescens</i>	NR, G5			X		23
Blue-winged Warbler	<i>Vermivora pinus</i>	S5, G5			X	WL, PIF	3
Bobolink	<i>Dolichonyx oryzivorus</i>	S5, G5			X	PIF	15
Brown Thrasher	<i>Toxostoma rufum</i>	S5, G5			X		21
Canada Warbler	<i>Wilsonia Canadensis</i>	S5, G5			X		13
Cooper's Hawk	<i>Accipiter cooperii</i>	S4, G5	SC		X		5
Eastern Meadowlark	<i>Sturnella magna</i>	S5, G5			X		14
Golden-winged Warbler	<i>Vermivora chrysoptera</i>	S4, G4	SC		X	WL, PIF	1
Louisiana Waterthrush	<i>Seiurus motacilla</i>	NR, G5			X		18
Northern Bobwhite	<i>Colinus virginianus</i>	S4, G5	G		X		1
Northern Goshawk	<i>Accipiter gentiles</i>	S4B, S3N, G4	SC		X		7
Olive-sided Flycatcher	<i>Contopus cooperi</i>	S5, G5			X	WL, PIF	3

Prairie Warbler	Dendroica discolor	S5, G5			X		12
Red-headed Woodpecker	Melanerpes erythrocephalus	S4, G5	SC		X	WL, PIF	1
Ruffed Grouse	Bonasa umbellus	NR, G5	G		X		20
Scarlet Tanager	Piranga olivacea	NR, G5			X		34
Sharp-shinned Hawk	Accipiter striatus	S4, G5	SC		X		9
Vesper Sparrow	Poocetes gramineus	S5, G5	SC		X		3
Whip-poor-will	Caprimulgus vociferous	S4, G5	SC		X	PIF	2
Willow Flycatcher	Empidonax traillii	S5, G5			X	WL, PIF	5
Wood Thrush	Hylocichla mustelina	S5, G5			X	WL, PIF	30
Worm-eating Warbler	Helmitheros vermivorus	S4, G5			X		1

Observed Rare Species and Significant Ecological Communities (source: NY Natural Heritage Program)


Rare plant and animals species with known populations within the watershed and documented examples of rare and high quality ecosystems within the watershed. Information regarding the locations of rare species is considered sensitive. The distribution of information which identifies the locations of rare species or their habitats may lead to the collection or disturbance of the animals and plants at those locations.

Key: TNC Status: For State and Global Rank explanations see www.natureserve.com ; Legal Status: FE = Federal Endangered; FT = Federal Threatened; SE = State Endangered; ST = State Threatened; G = Game species; SC = State Special Concern; NYNHP Species: Rare species tracked by the NY Natural Heritage Program; NYSDEC SGCN: Species of Greatest Conservation Need included in State Wildlife Plan; Other Ranking: Listed Partners in Flight WatchList					
Rare Birds		TNC Status	Legal Status	NYNHP Species	NYSDEC SGCN
Bald eagle	Haliaeetus leucocephalus	S2S3B, S2N, G5	ST, FT	X	X
Bicknell's Thrush	Catharus bicknelli	S2S3B, G4	SC	X	X
Rare Plants		TNC Status	Legal Status	NYNHP Species	NYSDEC SGCN
Blunt-lobe Grape Fern	Botrychium oneidense	S2S3, G4Q	SE	X	
Climbing Fern	Lygodium palmatum	S1, G4	SE	X	
Musk Root	Adoxa moschatellina	S1, G5	SE	X	
Rough Avens	Geum virginianum	S2, G5	SE	X	
Whorled Mountain-mint	Pycnanthemum verticillatum var. verticillatum	S1S2, G5T5	ST	X	
Rare Invertebrates		TNC Status	Legal Status	NYNHP Species	NYSDEC SGCN
Appalachian Tiger Beetle	Cicindela ancocisconensis	S1, G3	U	X	X
Natural Communities		TNC Status	Legal Status	NYNHP	NYSDEC SGCN
Hemlock-northern hardwood forest		S4, G4G5		X	
Beech-Maple Mesic Forest		S4, G4		X	
Spruce-Northern Hardwood Forest		S3S4, G3G4		X	
Mountain fir forest		S2, G3		X	
Mountain Spruce-Fir Forest		S2S3, G3		X	

References

- Allan, B.F., Keesing, F., and Ostfeld, R.S. 2003. Effect of Forest Fragmentation on Lyme Disease Risk. *Conservation Biology* 17(1): 267-272.
- Baldigo, B.P., Gallagher, E. Kelly, W., Warren, D.R., Miller, S.J., Davis, D., Baudanza, T.P., DeKoskie, D. and Buchanan, J.R. 2006. Restoring Geomorphic Stability and Biodiversity in Streams of the Catskill Mountains, New York, USA. *American Fisheries Society Symposium*, 2006.

- Curtis, P. D. (accessed 2004). Can QDM Improve Deer Management in Forested Landscapes? Department of Natural Resources, Cornell University. Ithaca, NY 14853
http://www.dnr.cornell.edu/ext/forestry/page/pubs/infobroch/by%20topic/quality_deer_management_curtis.htm.
- Evans, R.A. 2002. An Ecosystem Unraveling? Proceedings: Hemlock Woolly Adelgid in the Eastern United States Symposium, February 5-7, 2002, East Brunswick, New Jersey. <
http://www.fs.fed.us/na/morgantown/fhp/hwa/pub/proceedings/eco_unravel.pdf> (Accessed February, 2004).
- Kyker-Snowman, Thom. 2003. Rotten Logs and Sowbugs: the Role of dead wood. Massachusetts Association of Professional Foresters. <http://www.massforesters.org/coarse.htm> (Assessed 1/30/04).
- LoGiudice, K., Ostfeld, R.S., Schmidt, K.A., Keesing, F. 2003. The ecology of infectious disease: Effects of host diversity and community composition on Lyme disease risk. Proceedings of the National Academy of Sciences 100(2): 567-571.
- NYSDEC. 1993. Stocking Policy for the Greene County East Kill using 1990 Guidelines. New York State Department of Environmental Conservation, Stamford, NY.
- NY Natural Heritage Program Conservation Guides
 Online at: <http://www.acris.nynhp.org/>
- NYS Amphibian and Reptile Atlas
 Online at: <http://www.dec.state.ny.us/website/dfwmr/wildlife/herp/>
- NYS Breeding Bird Atlas
 Online at: <http://www.dec.state.ny.us/website/dfwmr/wildlife/bba/index.html>
- NYS Comprehensive Wildlife Conservation Strategy
 Online at: <http://www.dec.state.ny.us/website/dfwmr/swg/cwcsmainpg.html>
- NYS Threatened, Endangered & Special Concern Species List
 Online at: <http://www.dec.state.ny.us/website/dfwmr/wildlife/endspec/>
- Orwig, D.A. 2002. Stand Dynamics Associated with Chronic Hemlock Woolly Adelgid Infestations in Southern New England. Proceedings: Hemlock Woolly Adelgid in the Eastern United States Symposium, February 5-7, 2002, East Brunswick, New Jersey. <
<http://www.fs.fed.us/na/morgantown/fhp/hwa/pub/proceedings/dynamics.pdf>> (Accessed February, 2004).
- Pimentel, D., Wilson, C., McCullum, C., Huang, R., Dwen, P., Flack, J., Tran, Q., Saltman, T., Cliff, B. 1997. Economic and Environmental Benefits of Biodiversity. *BioScience* 47(11): 747-756.
- Smith, C.R., S.D. DeGloria, M.E. Richmond, S.K. Gregory, M. Laba, S.D. Smith, J.L. Braden, W.P. Brown, and E.A. Hill. 2001. An Application of Gap Analysis Procedures to Facilitate Planning for Biodiversity Conservation in the Hudson River Valley, Final Report, Part I: Gap Analysis of the Hudson River Valley and Part 2: Atlas of Predicted Ranges for Terrestrial Vertebrates in the Hudson River Valley. New York Cooperative Fish and Wildlife Research Unit, Department of Natural Resources, Cornell University, Ithaca, N.Y.
- Snyder, C.D., Young, J.A., Smith, D., Lemarie, D.P., and Smith, D.R. 2002. Influence of eastern hemlock (*Tsuga canadensis*) forests on aquatic invertebrate assemblages in headwater streams. *Canadian Journal of Fisheries and Aquatic Sciences* 59(2): 262-275.



Provided by NYSDEC, Hudson River Estuary Program
 Data Sources: NYS DEC Master Habitat Databank,
 NY Natural Heritage Program, February 2007

This map is informational only. This is not a regulatory map.

2.10 Recreational Opportunities

The Catskill Park is a mountainous region of public and private lands in Ulster, Greene, Delaware, and Sullivan counties. The natural and cultural heritage of the Catskill Region is inextricably linked to the unique high quality streams that course through its mountains and valleys and play a defining role in the character of its landscape. Recreation in and around these Catskill streams provides residents and visitors with a myriad of opportunities to reconnect with the natural world.

Catskill Forest Preserve

The East Kill watershed is surrounded by the steep mountains of the Catskills. The entire East Kill watershed is located within the Catskill Park “blue line” (Fig 2.10.1). Twenty-eight percent of the watershed is managed by the New York State Department of Environmental Conservation (DEC) as Wilderness and Wild Forest. The majority of this protected land is located at the creek’s headwaters and




Fig 2.10.1 East Kill Creek Watershed Parks and Preserves

lies within the Blackhead Range and Colgate Lake Wild Forests. Within these forests, approximately 2 miles (14%) of the East Kill flows through protected preserve. Colgate Lake, located in this area, is a popular tourist destination and is part of the East Kill’s headwaters.

In total, there are 6,315 acres of Forest Preserve and 11.8 miles of foot trails within the East Kill watershed (Figure 2.10.2). Black Head, Black Dome, and Thomas Cole Mountains make up the Blackhead Range Wild Forest peaks and are popular hiking

destinations. The East Kill watershed is surrounded by other preserves outside its borders including Windham High Peak Wild Forest to the north and Hunter Mountain Wild Forest and Indian Head Wilderness to the south. North/South Lake State Campground is located a few miles southeast of the watershed and Devils Tombstone State Campground is located a few miles directly south of the watershed. Information on hiking these and other locations, including trail descriptions, pictures, hiker reviews, topographic maps, and literature



View of the Blackhead Mountain Range (center-background) from Hunter Mountain summit

references can be found at www.localhikes.com. Hiking is permitted on all lands held by New York State. Hunting is allowed but subject to required NYS licenses and regulations.

The New York State Department of Environmental Conservation (DEC) manages lands in the forest preserve according to its classification in the Catskill Park State Land Master Plan (NYSDEC, 1985). Management recommendations are based on specific land characteristics and its capacity to withstand certain uses. These public uses include Wild Forest, Wilderness, Intensive Use and Administrative Use. DEC's *Catskill Forest Preserve Map and Guide* graphically depicts the locations of these different management areas and provides general background information about the Catskill Park and Preserve. This information can be obtained at DEC's regional offices. Locations of their offices are listed on DEC's website: <http://www.dec.state.ny.us/website/about/abtrull3.html>. There is also an interactive map on DEC's website called Environmental Navigator. This map shows the entire state of New York and can be magnified to specific locations, showing recreational attributes such as trails and parking facilities:

<http://www.dec.state.ny.us/cfm/extapps/statelands/index.cfm>.

Fishing


The East Kill is renowned for its fishing. Cold water temperatures and heavy forest cover especially favor trout populations. In 2005, it was stocked with 2,250 brown trout by

NYS DEC. Several fishing easements have been established along the East Kill near its confluence with the Schoharie Creek (Figure 2.10.3). Trout fishing season is open from April 1st through October 15th. A New York State Fishing License is required for those over 16 years of age. The basic state catch limit of five trout is applied to the East Kill and, as elsewhere, the practice of Catch and Release is voluntary, but encouraged. For more rules and regulations, or to view the fishing seasons for other fish species, visit DEC's Freshwater Fishing Regulations at:

<http://www.dec.state.ny.us/website/dfwmr/fish/fishregs/fishregsguide0608.pdf>.

References

NYSDEC. 1985. Catskill Park State Land Master Plan. New York State Department of Environmental Conservation (DEC), Albany, NY.




NYS DEC Forest Preserve Land

East Kill Watershed
 Fig 2.10.2

N

0 0.5 1 1.5 2 Miles

- Legend**
- BLACKHEAD RANGE WILD FOREST
 - COLGATE LAKE WILD FOREST
 - HUNTER MOUNTAIN WILD FOREST
 - KAATERSKILL WILD FOREST
 - WINDHAM HIGH PEAK WILD FOREST
 - PRIVATE INHOLDING
 - Lean-to
 - Trails
 - East Kill Watershed
 - Catskill Park




Fishing Easements
 (Bank designations are as you are looking downstream)

	Left
	Right

NYS DEC Public Fishing Easements

East Kill Basin

Fig 2.10.3



2.11 Stakeholders

Saturday, June 3, 2006, marked the official start to the stakeholder involvement portion of the Schoharie Creek and East Kill Stream Management Planning Projects. Greene County Soil and Water Conservation District (GCSWCD) and New York City Department of Environmental Protection (DEP) representatives presented information on the two agency's water resource programs and the various components of the stream management plans to the 100 participants. Presentations were followed by a question and answer session that included passionate stream management discussions that should lead to active participation in the planning process.



Schoharie Creek/East Kill Informational meeting, June 3, 2006.

Results from the June 3rd session reinforced that a critical component of the stream management planning process is public support and input for the project. To this end, the Schoharie/East Kill project team, and a professional consultant from the Consensus Building Institute, completed a survey of potential Project Advisory Committee (PAC) members to start gathering input for the management planning process (full results and recommendations below). Most interviewees described the stream as picturesque and aesthetically pleasing, historic and of great potential value. At the same time, it was also described as “out of control”, in need of attention, and unstable.

The survey information was further expanded upon at a facilitated workshop at which the following priority considerations for forming a project advisory committee to assist in the development and implementation of the stream management plans were decided upon: 1) representation of all key stakeholder groups; 2) manageable size (30 is probably about the biggest manageable group for full group discussions) 3) balance among stakeholder interests; 4) PAC membership should stay open through the planning process, so that new players can be added if appropriate. The group also decided the primary goals of the PAC should include: 1) building consensus among the key stakeholders on the goals, process and expected outcomes of the SMP process, based on broad public input; 2) engaging key stakeholders in reviewing information about the current conditions of the Schoharie Creek and East Kill and

its management, and gathering new information if necessary; 3) facilitating joint development of options for improving management, especially options that can contribute to multiple goals (environmental, economic, recreational and social); 4) prioritizing and integrating those options into a balanced management plan that says how resources will be allocated, who will do what and by when, and how to implement the preferred options; 5) mobilizing key stakeholders and the public to work together to generate the resources needed to implement the plan. Following the workshop a PAC was initiated that met several times prior to completion of this stream management plan. It should be noted that “completion” is a relative term. The stream management plans are designed to be updated with new assessments, knowledge or recommendations. Additionally, the PAC will develop annual action plans to prioritize recommendations. Therefore, the management planning process is not a stagnant process that concludes with the development of this stream management plan, but rather begins.

Occurring simultaneously with the Schoharie/East Kill watershed planning process was the Schoharie Turbidity Task Force. This project was designed to develop a turbidity reduction strategy for the Schoharie basin. The project included the surveying of stakeholder interests to better understand the challenges that turbidity poses to various interest groups (i.e. local residents, fishermen, water supply, local officials, highway crews, etc.). In addition, the hosting of a “turbidity summit” to present turbidity concerns within the Schoharie basin, possible best management practices to reduce turbidity and to gather input from ~ 100 attendees. Final turbidity reduction recommendations are scheduled to be completed in late 2007/early 2008.



Schoharie Turbidity summit break-out session (1 of 3), January 27, 2007.

All of the stakeholders listed in Table 2.11.1 have an interest in maintaining the East Kill as a well-functioning natural resource, and many of them have direct management influence over it. With the completion of the plan, the next phase will include review of the plan’s recommendations by the community, stakeholders and the Project Advisory

Committee. The plan will then be revised to ensure that it adequately reflect stakeholders’ concerns, and then presented to the various municipalities and agencies for formal adoption and implementation.

Table 2.11.1. Stakeholder groups within the Schoharie basin.	
Landowners	Federal Emergency Management Agency
Watershed Towns and Villages	Community Organizations
Greene County Soil and Water Conservation District	NYS Department of Environmental Conservation
Greene County Highway Department	USEPA
Army Corps of Engineers	Private utility companies
NYC Department of Environmental Protection	NYS Department of Transportation, including Region
NY State Emergency Management Office	Local Businesses

**Schoharie/East Kill Summary of Stakeholder Interests and Concerns
Final Report and Recommendations from the Consensus Building Institute**

Overview

The Consensus Building Institute (CBI) was asked by the DEP Stream Management Program, in cooperation with the GCSWCD, to help develop a Project Advisory Committee to provide public input and decision-making for the Schoharie/East Kill Stream Management Plans.

DEP and GCSWCD believe that in order to provide for maximum protection by multiple entities, developing a stream management plan must be a collaborative process among DEP, the local Soil and Water Conservation District(s), local governments, local nongovernmental agencies, watershed residents, and local business representatives. To promote collaboration, DEP and GCSWCD will create a Project Advisory Committee (PAC) of key stakeholders to help develop, and eventually implement, the Schoharie Stream Management Plan.

In order to develop an effective community engagement process that might result in diverse constituencies with ownership in the final plan, CBI was asked to conduct an impartial assessment of the interests and concerns of a broad range of stakeholders regarding the Schoharie and East Kill Watersheds. This assessment was presented at, and supplemented by, a focus group meeting of stakeholders on November 1, 2006.

The Consensus Building Institute is a not-for-profit [501 c(3)] organization based in

Cambridge, Massachusetts. CBI provides facilitation and mediation services to help public, private, and non-governmental organizations throughout the United States and internationally reach agreement on complex public policy matters.

Background

In response to damage of private property and public infrastructure caused by flood events, as well as ongoing water quality and fish habitat concerns, the GCSWCD, in cooperation with the DEP Stream Management Program, initiated the development of stream management plans for the Schoharie Creek and the East Kill.

The Schoharie–East Kill Watershed project is a continuation of a 10-year collaborative effort between the GCSWCD and DEP to promote sound stream management in the Catskills. The project involves conducting a thorough assessment of both streams, followed by the development of a detailed stream management plan for both waterways. While the project involves a significant number of activities, there are three main objectives: Complete a detailed assessment of the stream corridor to develop a reasonable, science based, understanding of the status and condition of the stream systems, including identification of stability problems; use data collected, in conjunction with watershed stakeholder discussions, to develop a detailed stream management plan for the watershed; conduct a demonstration project that will present a “hands-on” opportunity to restore stability at a selected site. Once completed, the Stream Management Plans (SMP) will provide an effective tool for government, citizens and other interested parties to manage the streams in a manner that will protect water quality, private & public property as well as the fisheries resource.

The Schoharie Creek main stem watershed drains large sections of Prattsville, Lexington, Jewett, Hunter, and the Villages of Hunter and Tannersville, and smaller sections of Ashland in Greene County. The East Kill watershed drains primarily the town of Jewett. Both the Schoharie Creek and East Kill are very “active” streams, noted for their tendency to quickly change from gentle waterways to raging streams under flood conditions. Damage from floods has been a way of life along these waterways for as long as humans have inhabited the watershed.

Methodology

The Consensus Building Institute conducted confidential, voluntary interviews with 9 individuals between August and October, 2006. In the interviews, CBI asked stakeholders questions about: perceptions and uses of Schoharie Creek and East Kill; interests, needs, and concerns associated with the Stream Management Plan, and; how and in what ways they would like to be involved in the watershed planning process. Interviews were done with individuals who fit into key stakeholder categories, including local government officials, local administrators, landowners, recreation groups, and federal agencies.

CBI staff used an established interview protocol as a general guide for conducting the interviews. The interviewer followed the general structure of the protocol, while allowing each conversation to follow the interests and comments of each interviewee. This summary is based on the views and opinions of interviewees. In addition, draft findings from these interviews were shared at a focus group meeting on November 1, 2006, with 25 attendees. Participants were asked to respond and add to these findings – these responses and additions are integrated into this summary.

Please note that the CBI's role is to provide accurate, impartial analysis of the situation in order to assist the DEP and GCSWCD in making recommendations on how to best proceed with a process able to engage the community in appropriate participation in the Stream Management Plan process and implementation. We are not an advocate for any particular outcomes or interests and are bound to conduct our work in a fair, deliberate, and non-partisan fashion. CBI staff is bound by the Association for Conflict Resolution (ACR) Code of Ethics: "The neutral must maintain impartiality toward all parties, maintain freedom from favoritism or bias either by word or by action, and commit to serve all parties as opposed to a single party". Please also note that the assessment is not a legal document, technical report or planning study, nor an exhaustive study of all the concerns of individuals and organizations with a stake in the Schoharie/East Kill Stream Management Plan. This final report is limited by the information gathered in the interviews and focus groups conducted and our interpretation of that information. Any errors or omissions are the sole responsibility of CBI.

Findings

Perceptions and Uses of Schoharie/East Kill Creek

Most interviewees described the stream as picturesque and aesthetically pleasing, historic and of great potential value. At the same time, it was also described as out of control, in need of attention, and unstable. They told stories of historic uses for the creek including swimming, boating, tubing, and fishing, but most said that these activities are quite curtailed in current times, due to the shallowness of the water. Others raised questions about the accuracy of the memory of a historically deeper stream, and questioned whether kayaking or tubing were ever popular on the stream. It was referred to as a great fishing stream, with consistent fishing and good retention of the stocked fish. It was also mentioned that there has been a reduction in insect and minnow life over the years. In addition to its beauty, the creek's value was described as an economic resource, an aquatic habitat, a source of water for the local ski area, conveyance for regulated water discharge from treatment plants, and a site for fishing.

Interests, Needs, and Concerns associated with the Stream Management Plan

Interviewees named a number of concerns they had about the stream. These included flooding, erosion, turbidity, backfill, invasive plants, storm water overflow, impacts of rapid development (especially potential future development), public access, landowner stewardship, and aquatic preservation.

Some comments from stakeholders include:

- Single biggest concern is that it needs to be controlled within its banks
- Flooding is getting worse. Extensive damage to town roads and personal property
- Within the town of Hunter, the community could use enhanced access
- Concerned about the potential impacts of new development, especially unplanned
- Need to enhance landowner stewardship and local commitment
- Additional development –especially unplanned – could increase storm water run off, sediment and turbidity
- Erosion has to be curtailed where clay banks are becoming exposed
- Turbidity exists, but the stream is one of the quickest in the area to clear up after

rain events

In speaking about their interests for the stream, stakeholders mentioned stream bank stabilization, studies of the stream beyond turbidity, including aquatics, restrictions on development, greater investment in structural improvements and preservation efforts, and a desire to see it restored back to a depth that would be useful for fishing, kayaking, tubing and swimming. There was also a call to avoid channelization of the stream, plan for and document the more-than-occasional flooding, and to identify and implement aquatic habitat improvement projects. Conversation arose about the history of gravel mining in the stream by local landowners, which is now prohibited, and is seen by many to have caused longer term instability. The group raised the question of how to meet objectives of flood mitigation & fish habitat improvement without gravel mining.

Interviewees had some ideas about how to bring about improvements in the stream, which could be included as part of the Stream Management Plan recommendations. These included requests to dig out the stream, eradicate non-native plants and educate people not to plant them, protect the natural features that help confine the flooding, and allow landowners to do work in the stream. Interviewees were also interested in strategies to prevent filling of banks, erosion, and flooding, improving the shading of the stream to reduce high August water temperatures that harm aquatic life, and a desire for new construction of berms and bridges to redirect the flow when it gets high. There were suggestions that the plan ensure that all initiatives were first the work of the local users, and that it raise awareness and interest of stakeholders, making the creek something important to them that they want to improve. This could be done through an offer of incentives and resources to towns and/or landowners to improve the stream. The idea was also raised of setting up volunteer opportunities – cleaning up streams, plantings, monitoring, pulling out Japanese Knotweed – for groups that might be willing/able to participate.

A focus group participant also mentioned that it would be wise to reduce the acreage requirement for disturbances that require stormwater management plans from the current requirement of over 1 acre because smaller disturbances may still impact water quality. Another attendee offered that reducing this limit would be too cumbersome on landowners and mentioned landowners could sue the water quality/quantity offender instead.

Recommendations for Local Involvement in the SMP process

Based on input from interviews and the focus group, as well as interests from DEP and GCSWCD in ensuring collective buy-in and ownership of the SMP recommendations, convening parties should establish a Project Advisory Committee. The most important considerations in deciding PAC membership should be 1) representation of all key stakeholder groups; 2) manageable size (30 is probably about the biggest manageable group for full group discussions); 3) balance among stakeholder interests (e.g. if there are 8 people from the Town of Hunter who want to participate, but only 1-2 people from other affected areas, the Hunter group should probably be asked to talk together to figure out which 2 or 3 can best represent their shared interests at any given meeting). PAC membership should stay open through the planning process, so that new players can be added if appropriate.

Given this, PAC members should include:

- Representatives of all Towns/Villages in these watersheds including: town supervisors, planning board members, highway superintendents and code enforcement officers;
- Any and all interested streamside landowners, with special efforts made to reach out to non-resident landowners because they do not elect Town officials and may not be represented by them;
- Any and all interested local businesses with a stake in the streams;
- Representatives from state and federal government agencies;
- Representatives from non-governmental organizations and recreation groups.

The primary goals and tasks for the group should include:

- Building consensus among the key stakeholders on the goals, process and expected outcomes of the SMP process, based on broad public input;
- Engaging key stakeholders in reviewing information about the current conditions of the Schoharie Creek and East Kill and its management, and gathering new information if necessary;
- Facilitating joint development of options for improving the management of the Schoharie and East Kill, especially options that can contribute to multiple goals (environmental, economic, recreational and social);

- Prioritizing and integrating those options into a balanced management plan which says how resources will be allocated, who will do what, by when and how to implement the preferred management options;
- Mobilizing key stakeholders and the public to work together to generate the resources needed to implement the plan.

The PAC should ensure inclusion, participation, and a real voice of members in decision making, and should seek to hold participants accountable for collecting and offering perspectives from those they represent. Given time constraints, the PAC should meet at least 2-3 times before April, 2007, and should then commit to working together to adopt and implement the recommendations of the SMP.

The PAC meetings should include a 2-way exchange of information. They are an opportunity for the project team to share their progress and current thinking about the SMP and its recommendations, to explore outstanding questions or disagreements among the group (such as “how deep is the stream? How deep did it used to be?”) and also for the PAC members to build consensus on SMP recommendations and priorities for funding.

In addition to the PAC, GCSWCD should offer other opportunities for the participation of the public at large. Those opportunities might include:

- Hosting 1-2 public meetings together with the PAC, to share information about progress so far and draft ideas on recommendations and priorities, to provide an opportunity for community input;
- Creating a website and/or regular electronic and print updates which the public can use to follow the progress of the SMP;
- Involving them in data gathering activities, e.g. putting monitoring devices on their property or responding to questionnaires;
- Providing input to members of the PAC, and/or attending meetings of the PAC as observers.

By creating opportunities like these, NYC DEP and GCSWCD can broaden and deepen stakeholder engagement in the SMP process.

2.12 Agency Contacts and Program Resources

Technical Assistance

A wealth of information and assistance is available to local municipalities, landowners, and businesses in the Catskill/Delaware watershed. Services are wide ranging through a variety of programs. Although funding and grant opportunities may not always be a possibility, the organizations listed below offer a variety of solutions for water quality, infrastructure, and property protection. Please do not hesitate to contact these resources with questions and requests. Many of these organizations also offer grant and other funding opportunities. Please see the grant resources list at the end of this section for more information on monetary support (Table 2.12.1).

Soil & Water Conservation Districts (SWCD)

With a conservation district in each county, these local entities provide a variety of services to its local constituency. Most districts focus on offering agricultural assistance with best management practices (BMPs) through design, installation, and oversight. These BMPs include water management such as diversions, barnyard management systems, manure storages, grazing systems, and animal water systems. Other services include riverfront revitalization, plant materials supply, environmental education, permit assistance, flood mitigation, and stream restoration. The SWCDs are often a good starting place for information and assistance. If they cannot help you, they can most likely point you in the right direction.

Greene: Executive Director

(518) 622-3620

Schoharie: Executive Director

(518) 234-4092

Delaware: Executive Director

(607) 865-7161/7090

New York City Department of Environmental Protection (NYC DEP)

<http://www.nyc.gov/html/dep/html/watersup.html>

The Bureau of Water Supply works closely with landowners to achieve goals in an environmentally sensitive manner. NYC DEP has a variety of programs that assist

landowners with the management of their property and streams. Please see below for a brief description of the various programs.

Land Acquisition: In 1997, the DEC issued a permit that allowed the DEP to acquire land for the purpose of watershed protection. The acquisition of land is one of the best ways to ensure the ongoing prevention of pollution and to prevent future water quality problems from occurring as a result of adverse development close to critical natural features and reservoir intakes. Purchase of land at fair market value or placement in an easement is negotiated only from willing sellers. Interested parties should contact the Land Acquisition Program @ 1-800-575-LAND or (845) 340-7540.

Stream Management: DEP's Stream Management Program was established in 1992, and formalized in 1996, as one of the watershed community partnership programs included in the 1996 Watershed Memorandum of Agreement. Its mission is to establish long-term stewardship of streams through a watershed-scale, community-based, geomorphic approach, and the development of Stream Management Plans for priority sub-basins in the NYC Water Supply West-of-Hudson (WOH) watersheds. Essential to achieving this goal is the provision of technical assistance to local municipalities, landowners, and businesses within the watershed. Staff members also offer training and educational programs regarding these topics. Concerns or requests for service, should be made to the Stream Management Program at (845) 340-7628.

Land Management: The Land Management Program develops land resource management plans for DEP properties, conducts a recreational review, and develops basin plans, incorporating specific property by property uses and stewardship. In addition, the DEP implemented a public access program that of August, 2006 had made 65% of acquired lands in the Schoharie basin available for recreational purposes like hiking, hunting, and fishing. For additional information call (845) 340-7541.

The DEP also oversees a number of other programs like the watershed agricultural and watershed forestry programs, sewer and septic maintenance, economic development, and watershed education through the Catskill Watershed Corporation (CWC). Please see the CWC description below for more details.

New York State Department of Environmental Conservation (NYS DEC)

www.dec.state.ny.us

Many water related programs are offered by the NYS DEC. The agency has various divisions, which handle watershed assessment and management, environmental education, fisheries, and flood protection. Information about the DEC stocking schedule, fishing licensing, and access points is available at

<http://www.dec.state.ny.us/website/dfwmr/fish/index.html> or by calling (607) 652-7366 for Region 4.

To receive information regarding any flooding issues and the National Flood Insurance Program, see <http://www.dec.state.ny.us/website/dow/bfp/gisfpm/index.htm> or call (518) 402-8141 about flood control projects or (518) 402-8146 about flood plain management.

In addition to the above services, the DEC is also the regulatory agency for the state of New York's waterways. Having classified Catskill streams, the DEC requires a Protection of Waters Permit for disturbing the bed or banks of a stream. Please contact the following for direction and advice.

Greene/Delaware/Schoharie Bureau of Habitat

65561 St Hwy 10

Stamford, NY 12167

(607) 652-2645

U.S. Army Corps of Engineers (ACOE) New York District

www.nan.usace.army.mil/index.htm

The Army Corps of Engineers has a variety of duties related to stream management. If a municipality or landowner wishes to install a water-related structure, dredge or fill a stream, or affect a wetland area, ACOE will often assign a field technician to visit the site in order to evaluate the need for a federal permit. ACOE also offers engineering designs and other technical expertise. In addition, they are available for planning, designing, and

constructing flood control projects. For a field technician contact the appropriate office listed below:

Delaware/Greene/Schoharie: (518) 273-7420

Catskill Watershed Corporation (CWC)

www.cwconline.org

The CWC is a not-for-profit corporation with a dual goal: to protect the water resources of the New York City Watershed west of the Hudson River, while preserving and strengthening communities located in the region. Although the CWC is mainly a source of funding (see grant information section), they can also provide technical assistance. Pertinent programs for Catskill/Delaware stream stakeholders include the Stormwater Controls for New Construction, Stormwater Retrofit, Septic System Rehabilitation and Replacement, and Alternate Design Septic Program. For more information call (845) 586-1400.

The Septic Rehab and Replacement program, administered by The Catskill Watershed Corporation (CWC) reimburses permanent residents 100 percent of eligible costs of repairing or replacing a failed septic system. Non-primary residents are reimbursed 60 percent of eligible costs. Eligible systems must be a one or two family residence or home-business combination using less than 1,000 gallons per day, and be located in the NYC Watershed in Delaware, Greene, Schoharie, Sullivan or Ulster Counties. The septic system must be located within 150-feet of a watercourse or within 500 feet of a reservoir or reservoir stem in the West-of-Hudson (WOH) Watershed. This program does not pay for new septic systems serving newly constructed home; or for new or repaired systems intended for commercial or institutional use (CWC, 2006).

The Septic Maintenance program is intended to extend the life of septic systems serving one and two family households in the West-of-Hudson Watershed. This program is open to homeowners anywhere in the NYC WOH watershed, who have had a new, or replacement septic system installed after January 21, 1997, and at least three years ago. This program reimbursed homeowners for up to 50% of the eligible cost for septic system inspections and pump-outs.

Watershed Agricultural Council (WAC)

www.nycwatershed.org

WAC offers the Watershed Agricultural Program and the Watershed Forestry Program. WAC subcontracts with local, state, and federal agricultural assistance agencies, Cornell University, and the private sector to provide planning, education, training, engineering, scientific, and administrative support.

Watershed Agricultural Program (WAP)

WAP strives to protect the high water quality from agricultural nonpoint source pollution through the planning and implementation of Best Management Practices (BMPs) on farms. Using traditional and non-traditional BMPs, WAP strives to offer a variety of alternatives to farmers that promote the health of their land and the stream. Some specific programs are Whole Farm Planning, the Conservation Reserve Enhancement Program, Nutrient Management Planning, and Small Farm Program. Call (607) 865-7790 or email info@nycwatershed.org with questions or requests.

Conservation Reserve Enhancement Program (CREP)

This program is available to current agricultural landowners or landowners who may not currently farm land, but whose property has a history of agricultural use. CREP is a program for promoting the health of streamside vegetation by providing rental payments for buffer lands that are taken out of production, as well as 100% funding for tree/shrub planting. This program also helps landowners implement stream fencing and livestock watering facilities and other BMPs.

Watershed Forestry Program (WFP)

The Watershed Forestry Program is a voluntary partnership between New York City and the upstate forestry community that maintains well-managed forests as a preferred land use for watershed protection. In 2001, forests covered approximately 85% of the Schoharie basin land area, and a majority of this forest land is privately owned and managed by thousands of individual landowners. To promote forest stewardship and encourage long-term investment in private forestry, the Forestry Program offers cost-sharing to landowners for developing 10-year forest management plans written by qualified professional foresters.

Participating landowners must own at least 10-acres of forest land in the watershed. The Forestry Program also offers a variety of cost-sharing, technical assistance and other incentive programs to both loggers and landowners for implementing certain forestry practices that protect water quality, such as properly installing new timber harvest roads and stream crossings or remediation of existing forest roads that have documented erosion problems. Owning a watershed forest management plan is actually a prerequisite for many of these programs. Forest landowners may also attend a variety of educational workshops and other training events that are periodically sponsored throughout the watershed. For more information, call (607) 865-7790 or email forest@catskill.net.

National Rural Water Association

www.nrwa.org

The National Rural Water Association is a non-profit federation of State Rural Water Associations. Their mission is to provide support services to State Associations who have more than 22,000 water and wastewater systems as members. See description below for New York state contact information.

New York Rural Water Association

http://www.nyruralwater.org/technical_assistance/technical.cfm

New York Rural Water Association (NYRWA) is a not-for-profit group organized in 1979 with the goal of promoting the development, improvement, and sound operation of rural drinking water and wastewater systems throughout New York State. New York Rural Water Association recently expanded its scope to offer training, technical, and administrative assistance to rural communities on solid waste management matters as well. Contact (518) 828-3155, or visit nyruralwater.org.

Federal Emergency Management Agency (FEMA)

<http://www.fema.gov/>

www.msc.fema.gov

FEMA is the federal government agency responsible for administering emergency and disaster relief, recovery, planning and preparedness programs across the United States and territories. While FEMA's most apparent role is emergency response and recovery, its role in risk reduction through the establishment of building codes and administration of

insurance programs, like the national flood insurance program, provide protection against losses of life and property in the case of an emergency or natural disaster. Based in Washington, FEMA operates regional offices across the United States including the Region II office in New York City, covering New York State. FEMA works in cooperation with other federal agencies and State and local emergency response entities such as the State Emergency Management Office (NYS SEMO) and county Emergency Management officials (please see below). FEMA provides training to state and local officials on most aspects of their work including emergency response, disaster response planning, hazard mitigation planning, code interpretation and enforcement. Following a Presidentially declared disaster, FEMA's assistance can be available to state and local government, private individuals, and businesses.

Floods are the most common disaster that would require FEMA involvement with Catskill watershed communities. To protect against flood damages and the loss of life associated with flood events, FEMA provides the following types of assistance:

- Administration of the National Flood Insurance Program (NFIP). Through this program FEMA prepares flood insurance rate maps (FIRMs) that define where floodwaters are likely to cause damage to property. These maps provide communities with a tool to prevent losses through the limitation of building and flood plain modification within these flood zones (Maps provided at FEMA's Map Service Center where you can access the most current FIRM maps: www.msc.fema.gov).
- Management of hazard mitigation programs that help communities identify and modify situations and places at risk during flood events. This would include the development of hazard mitigation plans prepared by communities to help the community reduce or avoid threats to life or property during flood events.
- Following flood events that are declared by the President to be a disaster for a specific county, FEMA typically provides assistance for temporary housing, clean-up, repairs to private structures and repairs to public infrastructure. The availability of this assistance depends on the magnitude of the disaster and the types of losses incurred by the county and its residents. The Small Business Administration also can provide assistance with low interest loans to private business. FEMA programs are

modified frequently and therefore the type and level of assistance will vary from event to event.

- FEMA plays its most important role as a coordinator of response and information in times of a disaster.

To contact the FEMA Region II office, please call (212) 680-3600.

New York State Emergency Management Office (NYS SEMO)

www.semo.state.ny.us

As stated above, the New York State Emergency Management Office is the state entity for pre- and post disaster assistance. Like FEMA, the state office provides planning and resources through cooperation with local governments, volunteer organizations like Red Cross, and the private sector. Where FEMA is primarily involved immediately after a disaster event, SEMO provides long-term recovery solutions. The state agency is more involved in the day to day planning and preparation for disaster response. Below are summaries of some of SEMO's major programs.

Mitigation: This may be one of SEMO's most influential programs by providing preventative assistance to communities within the Catskills. Mitigation efforts intend to reduce negative impacts of floods and other major disasters by preparing pre-disaster planning. This program also aims to identify potential threats and repeatedly damaged structures and to offer positive solutions to reduce future losses and protect against the loss of life and property. It is the intention that preventative efforts will greatly reduce the cost of recovery and will also reduce the loss of property. SEMO manages a Hazard Mitigation Grant program available to communities that prepare hazard mitigation plans. Communities preparing the plan are eligible for grant program funds to implement hazard mitigation projects following Presidentially declared disasters within New York State. Individuals living in communities with plans may benefit from the program through the reduction in flood insurance rates.

Disaster Recovery Assistance: Recognizing that not all disasters can be prevented, this program aims to provide local assistance for faster recovery by coordinating public

assistance funds, disaster housing assistance, individual family grants, and small business administration assistance.

Other Emergency Assistance: SEMO also provides a variety of services during times of emergency. These services include state of the art communications, information dissemination, and emergency operation coordination. Call the Emergency Coordination Center at (518) 929-2200 with questions or requests.

Cornell Cooperative Extension (CCE)

www.cce.cornell.edu

Cooperative Extension builds partnerships and coalitions with individuals, communities, organizations, government agencies, and businesses around issues of mutual concern; develops local leaders who use CCE knowledge to inform decisions; promotes youth development through 4-H clubs and other experiences; strives to help participants make informed choices using the best knowledge available; connects learners with educational resources found in locations throughout the world; consults with individuals and groups on multiple topics; provides resources via technologies such as the World Wide Web, satellite, and compressed video.

Greene: (518) 622-9820

greene@cornell.edu

Delaware: (607) 865-6531

delaware@cornell.edu

Schoharie: (518) 234-4303

schoharie@cornell.edu

The Greene County CCE Agroforestry Resource Center

<http://arc.cce.cornell.edu/>

The Agroforestry Resource Center provides educational programs, supports research, and promotes collaboration among organizations concerned with sustaining the forest dominated landscape of the Catskill Mountain/Hudson Valley Region. Agroforestry is the art and science of integrating timber production with other crops and forest related income opportunities. The goal is to promote sustainable practices that will provide economic benefits to landowners and communities while preserving forested areas so they continue to

provide ecological, economic and public health benefits to the surrounding human population.

Natural Resources Conservation Service (NRCS)

www.nrcs.usda.gov

NRCS puts nearly 70 years of experience to work in assisting owners of America's private land with conserving their soil, water, and other natural resources. Local, state and federal agencies and policymakers also rely on their expertise. They deliver technical assistance based on sound science and suited to a customer's specific needs. Cost shares and financial incentives are available in some cases. Most work is done with local partners. NRCS's partnership with local conservation districts serves almost every county in the nation, and the Caribbean and Pacific Basin. Participation in our programs is voluntary. Please see below for local contact information.

<i>Greene:</i>	Ghent Service Center (518) 828-4385	<i>Schoharie:</i>	Cobleskill Service Center (518) 234-4377
<i>Delaware:</i>	Walton Service Center (607) 865-4005		

United States Geological Survey (USGS)

ny.water.usgs.gov

The USGS provides the Nation with reliable information about the Earth to minimize the loss of lives and property from natural disasters, to manage biological, water, mineral, and energy resources, to enhance and protect the quality of life, and to contribute to wise economic and physical development. The USGS provides a variety of assistance related to the four main categories of biology, geography, geology, and water. The water division is broken down into ground water, surface water, and water quality. Individuals can find a multitude of data throughout the website, search various resource databases, and view a number of maps. For more information call the Troy office at (518) 285-5600.

Catskill Forest Association (CFA)

www.catskillforest.org

The Catskill Forest Association is a non-profit organization dedicated to enhancing all aspects of the forest in New York's Catskill region. CFA offers educational programs at all levels, from one-on-one site visits at landowner properties to group woods-walks, workshops and seminars. School-based activities include classroom visits and teacher training such as the Watershed Forestry Institute. CFA is also active in advocating for proper forest management, as well as promoting the economic development of viable markets for a variety of forest products. For more information, email cfa@catskill.net or call (845) 586-3054.

Catskill Center for Conservation and Development (CCCD)

www.catskillcenter.org

The Catskill Center is a non-profit organization working to protect the cultural, historic, and natural resources of the Catskill Mountains. The CCCD has a few integrated program areas:

Land Conservation & Natural Resource Protection: This program identifies, monitors, and engages in effective actions to protect and preserve sensitive, ecologically significant, aesthetically, or recreationally critical lands and waters.

Community Outreach and Planning Assistance: This program provides technical support to rural communities in the Catskills on grant-writing, planning, land use, zoning, subdivision, community empowerment, main street revitalization, regional forums, conferences and workshops, producing reports and publications, and public policy development.

Education: This program consists of a curriculum entitled The Catskills: A Sense of Place, which is a series of five modules on the water resources, geography and geology, ecosystems, human history, and culture and arts of the Catskills. A Sense of Place is designed to give children a better awareness, understanding, and appreciation of the distinctive features of our area. In addition, The Center has partnered with Hudson Basin River Watch to support advanced water quality monitoring efforts by adult volunteer groups. Lastly, they

host a hike, lecture, and recreation series for our membership and the general public throughout the year. Visit their website at catskillcenter.org or call (845) 586-2611.

Trout Unlimited (TU)

www.tu.org

Trout Unlimited's mission is to conserve, protect and restore North America's trout and salmon fisheries and their watersheds. TU accomplishes this mission on local, state and national levels with an extensive and dedicated volunteer network. Local TU members have been active in many aspects of stream management planning throughout the Catskill/Delaware watershed. Not only do they participate in public meetings, legislative activities, and volunteer events, but TU has also funded research projects such as the "Economic Impact Assessment of the Beaverkill-Willowemoc Trout Fishery" to promote improved trout habitats and stream health. Please contact the following local chapters for further information:

Ashokan-Pepacton 559: (845) 657-2312

Catskill Mountain 028: (845) 334-9323

Columbia Greene RVW 569: (518) 851-9442

Greene Land Trust

<http://www.greenelandtrust.org/>

Recognizing the need for an organizational structure that would insure long-term protection and management of critical habitat lands, the Greene County Soil & Water Conservation District and Greene County Industrial Development Agency funded the establishment of the Upper Hudson—Northern Catskill Natural Resource Trust, which later filed with the NYS Dept of State to operate under the assumed name of Greene Land Trust. While the GLT was founded to assist the habitat protection efforts, it is an autonomous organization that continues to grow and address a range of local natural resource issues in the Greene County area. Contact Rene Vanschaack at (518) 622-3620 or info@greenelandtrust.org.

ESRI Environmental Conservation Program (CSP)

www.conservationgis.org/aaesrigrants.html

This program provides donations and discounts of GIS software, data, books, and training. It offers free on-line live workshops. The overall goal of the ECP is to support conservation groups in acquiring, learning, and using GIS tools and methods. ECP has a particular focus on appropriate levels of technology for locally sustainable programs. Its goal is not to throw out one-off donations into a vacuum with no forethought, but to build permanent, locally based support structures that provide ongoing evolutionary growth in GIS skills. Email ecp@esri.com for detailed information.

Table 2.12.1. Funding Sources and Agency Contacts

Name	Focus	Due Date	Contact	Award Example	Notes/Priority	on-the-research	Planning	> \$20K	\$20K to < \$100K	Range
Natural Resources Conservation Service										
Conservation on Private Lands http://www.nfwf.org/AM/Template.cfm?Section=Browse_All_Programs&CONTENTID=3971&TEMPLATE=/CM/ContentDisplay.cfm	Projects that engage private landowners, primarily farmer and ranchers, on the ground project.		Eastern Partnership Office, Director, Peter Stangel 404-679-7099		Partnership with NRCS or local conservation districts, priority given to landscape, watershed scale projects integrating agriculture, forestry, and ranching that benefit fish and wildlife.	X		X	X	10K-150K
Emergency Watershed Protection http://www.nrcs.usda.gov/programs/ewp/factsheet.html	Projects support such work as clearing debris from clogged waterways, restoring vegetation, and stabilizing river banks.	on-going	Highland Service Center 845-883-7162		The measures that are taken must be environmentally and economically sound and generally benefit more than one property owner.	X				
National Oceanic and Atmospheric Administration										
Community-based Restoration http://www.nmfs.noaa.gov/habitat/restoration/projects_programs/crp/index.html	Provides funds for small-scale locally driven habitat restoration projects that foster natural resources stewardship within communities.		Robin Bruckner robin.bruckner@noaa.gov 301-713-0174	Provides funding to implement on-the-ground habitat restoration projects to benefit marine, estuarine and riparian habitats.		X		X	X	14K-8 mil.
Federal Emergency Management Agency										
Flood Mitigation Assistance http://cfpub.epa.gov/fe/fund/list1.cfm?program=31	Program helps states and communities identify and implement measures to reduce or eliminate the long-term risk of flood damage to homes and other structures.	established by states	Public Assistance Branch, Recovery Division FEMA, DHS 500 C Street, SW Washington, DC 20472 202-646-4262		Two types offered: planning and project grants for National Flood Insurance Program (NFIP) participating communities.	X	X			
U.S. Fish and Wildlife Service										
North American Wetlands Conservation Act Grants http://www.fws.gov/bird/habitat/Grants/index.shtml	Standard and small grants programs help deliver funding to on-the-ground projects through the protection, restoration, or enhancement of an array of wetland habitats.		Standard-David Buie 703-358-2266 Small-Keith Morehouse 703-358-1888	See award examples at: http://birdhabitat.fws.gov/hawca/grants.htm			X	X	X	small=<75K standard=75K-36.2mil.
Partners for Fish and Wildlife http://ecos.fws.gov/partners/viewContent.do?viewPage=home	Restoring former and degraded wetlands, native grasslands, stream and riparian areas, and other habitats to conditions as natural as feasible.	on-going	Carl Schwartz 607-753-9699	The program has partnered landowners to restore wetlands, prairie grassland, and in-stream aquatic and riparian habitat.	Provides technical and financial assistance to landowners interested in voluntarily restoring or otherwise improving native fish & wildlife habitat on their lands.	X		X	X	<25K

Table 2.12.1.1. Funding Sources and Agency Contacts, Continued

Name	Focus	Due Date	Contact	Award Example	Notes/Priority	on-the-ground research	> \$20K	\$20K to < \$100K	Range
Cooperative State Research, Education, and Extension Service									
National Integrated Water Quality http://www.usa.waterquality.org/intergrated	Supports integrated research, education, and extension projects, as well as, extension/education projects to address water quality issues at the watershed scale.		Mike O'Neill National Program Leader, Water Quality 202-205-5952 moneill@csrees.usda.gov	See award examples at: http://www.usawaterquality.org/projects	Grant awards to be made to four-year degree granting institutions.	X		X	85K-1.3 ml.
State Emergency Management Office									
http://www.semo.state.ny.us	Provides leadership, planning, education, and resources to protect lives, property and the environment.	on-going	State Emergency Coordination Center 518-292-2200 postmaster@semo.state.ny.us			X	X	X	
Catskill Watershed Corporation									
Catskill Fund for the Future http://www.cwconline.org/programs/econ_dev/econ_dev.html	Funds will be used to make loans and grants to businesses and organizations proposing environmentally responsible projects.	rolling basis	Michael Triolo, Economic Development Director, triolo@cwconline.org Phil Sireci, sireci@cwconline.org	Delhi received money for establishment of Riverwalk Community Park (purchase of riparian property and development of a village riverfront area with canoe access.	This fund program includes a variety of grant and loan programs.	X	X	X	
Septic System Rehabilitation and Replacement http://www.cwconline.org/programs/septic/rehab.html	This program reimburses homeowners for repairing or replacing damaged septic tanks.		Leo LaBuda labuda@cwconline.org John Jacobson jjacobson@cwconline.org 845-586-1400	Limited to properties in the five-county West-of-Hudson Watershed whose septic systems or property centroids lie within 100 feet of a watercourse. 60% and 100% of eligible costs for non-primary and primary landowners, respectively.		X	X		60%-100% of costs

Table 2.12.1. Funding Sources and Agency Contacts, Continued

Name	Focus	Due Date	Contact	Award Example	Notes/Priority	on-the-ground research	> \$20K	\$20K to < \$100K	Range
Catskill Watershed Corporation continued									
Stormwater Controls for New Construction http://www.cwconline.org/programs/sfrm_wtr/sfrm_wtr.html	Program to design and construct runoff and erosion control measures.		Thomas De John Professional Engineer tdejohn@cwconline.org	One project will improve stormwater collection and treatment on Railroad Ave., a project intended to decrease pollution and nutrient loading and reduce flow to the village's wastewater treatment plant.		X		X	
Stormwater Retrofit http://www.cwconline.org/programs/sfrm_wtr/sfrm_wtr.html	Program to provide funds for stormwater management needed to correct or reduce existing erosion, polluted runoff or other problems associated with stormwater.		Thomas De John Professional Engineer tdejohn@cwconline.org		Projects to implement stormwater BMPs that reduce erosion and/or pollutant loading associate with conditions existing on/or before January 21, 1997 are eligible to apply.	X			
Public Education http://www.cwconline.org/programs/pub_edu/pe.html	Projects that would increase awareness of the region's environment, its natural beauty, and human history.		Diane Galusha galusha@cwconline.org	Tri-Valley Central School, Grahamsville, water monitoring equipment to expand the agricultural and environmental studies programs to include water quality examinations; Ernest Myer School, to bring Streamwatch.		X	X		\$750 to \$10K
Local Technical Assistance Program http://www.cwconline.org/programs/tech/tech.html	Program addresses projects that enhance pollution prevention management plans or regulations intended to reduce existing /potential erosion and/or pollutant loading or improve the vitality of watershed communities		Thomas De John Professional Engineer tdejohn@cwconline.org	New Program	Preference given to projects that involve municipalities with commitment to adopting management plans, local law, or study recommendations.		X	X	<\$50K

2.12.1.1. Funding Sources and Agency Contacts, Continued

Name	Focus	Due Date	Contact	Award Example	Notes/Priority	on-the-ground research	> \$20K	\$20K to < \$100K	Range
National Fish & Wildlife Foundation									
Bring Back the Natives http://www.nfwf.org/AM/Template.cfm?Section=Browse_All_Programs&TEMPLATE=/C/M/ContentDisplay.cfm&CONTENTID=4095	Funds on-the-ground efforts to restore native aquatic species to their historic range. Can involve riparian habitat restoration, moving streams toward stability and supporting native aquatic communities.	two-decision cycles/year	Corey Grace 415-778-0999 corey.grace@nfwf.org		Projects involving sensitive or listed aquatic species. Does not fund basic research or monitoring.	X		X	
Native Plant Conservation Initiative http://www.nfwf.org/AM/Template.cfm?Section=Stewardship&TEMPLATE=/CM/ContentDisplay.cfm&CONTENTID=3966	Projects that protect and enhance, and/or restore native plant communities on public and private land, including protection and restoration, information and education, and inventory and assessment.	two-decision cycles/year	Ellen Gabel 202-857-0166 ellen.gabel@nfwf.org		There is a strong preference for "on-the-ground" projects that involve local communities and citizen volunteers in the restoration of native plant communities. Projects that include a pollinator conservation component are also encouraged.	X			10K-50K
Five-Star Restoration Matching Grants Program http://www.nfwf.org/AM/Template.cfm?Section=Browse_All_Programs&TEMPLATE=/C/M/ContentDisplay.cfm&CONTENTID=4502	Supports community-based wetland, riparian, and coastal habitat restoration projects that build diverse partnerships and foster local natural resource stewardship through education, outreach and training activities.		Amanda Bassow 202-857-0166 amanda.bassow@nfwf.org	The Mahopac High School will create a half-mile of wetlands along SEAC Creek in front of Mahopac High School by re-contouring the stream corridor and planting native trees and shrubs. The project will serve as a local example of ecological restoration and will be designed as an outdoor classroom for student educational use.	Preference will be given to projects that: Are part of a larger watershed or community stewardship effort; include specific provisions for long-term management, monitoring, and protection; and demonstrate the value of innovative, collaborative approaches to restoring the nation's waters.	X			5K-20K
Watershed Agricultural Council									
NYC Watershed Forestry Program http://www.nycwatershed.org/	Provides cost-sharing incentives and technical assistance to watershed forest landowners to promote forest management planning and to help establish streamside buffers.	rolling assistance	Watershed Forestry Program 607-865-7790 1-800-662-1220 info@nycwatershed.org		Assistance from this program could be used to establish additional grants from matching programs that require existing challenge funds and partnerships.		X	X	

Table 2.12.1. Funding Sources and Agency Contacts, Continued

Name	Focus	Due Date	Contact	Award Example	Notes/Priority	on-the-ground	research	planning	> \$20K	\$20K to < \$100K	Range
The Conservation Fund											
Kodak American Greenways Award http://www.conservationfund.org/article=2372	Small grants to stimulate the planning and design of greenways in communities throughout America.	03-01 to 06-01 each year	American Greenways Program 703-525-6300 postmaster@conservationfund.org	North American Water Trails received grant money for its development, enjoyment and stewardship or recreational water trails.	Grants used for appropriate expenses needed to complete greenway project including planning, technical assistance, legal and other costs.		X	X			
ESRI Conservation Program											
Software/Training Donations	Provides donations and discounts of basic GIS software and books.	on-going	grant@esri.com		Does not provide hardware or cash.						
Conservation Program http://www.conservationgis.org/aesrigrants.html Resource Conservation District Grant http://www.esri.com/grants/esri/conservation.html	Provides donations and discounts of GIS software, data, books, and training.	on-going	scp@esri.com rcdgrant@esri.com		Does not provide hardware or cash.						
Tech Grants											
http://www.techfoundation.org	TechFoundation is committed to bringing financial resources, technology solutions and management expertise to non-profits to strengthen the social sector.		techgrants@techfoundation.org 617-354-7500	Colorado Environmental Coalition www.ourcolorado.org	Awardees selected for focus on projects that will bring quality technology resources.	X	X	X			
Earthwatch Institute											
Research Program http://www.earthwatch.org/research	Supports field research worldwide in biological, physical, social, and cultural sciences.	on-going	Earthwatch Institute 1-800-776-0188 info@earthwatch.org	Monitor water quality. Inventory, monitoring, or restoration of watershed environments.	Grants cover cost of maintaining volunteers and research field staff not PI salaries, capital equipment, or overhead.			X	X	X	
Toshiba America Foundation											
www.toshiba.com/taf/apply.html	Contribute to projects designed by classroom teachers to improve science and math education.	on-going	212-596-0620 foundation@tai.toshiba.com	MS received grant for earth science students to conduct a water quality study in their area.		X		X	X		

2.13 Stream-related Activities and Permit Requirements

NYS DEC Permit Requirements

Certain kinds of human activities can have a detrimental impact on water resources. The policy of New York State is to preserve and protect lakes, ponds, rivers and streams, as set forth in the Environmental Conservation Law (ECL) Title 5 of Article 15. To implement this policy, the New York State Department of Environmental Conservation created the Protection of Waters Regulatory Program.

All waters of the State have a classification and standard designation based on existing or expected best usage of each water or waterway segment. The classification AA or A is assigned to waters used as a source of drinking water. Classification B indicates a best usage for swimming and other contact recreation. Classification C is for waters supporting fisheries and suitable for non-contact activities. Classification D, the lowest classification standard, reflects a best usage for fishing.

Waters with classifications, A, B, and C may also have a standard of (t), indicating that it is able to support a trout population, or (ts) indicating that it supports trout spawning. Special requirements apply to sustain these waters that support these valuable and sensitive fisheries resources. The East Kill and most of its tributaries have a classification and standard of C(t) or higher, and as such are subject to the stream protection provision of the Protection of Waters regulation.

A Protection of Waters Permit is required for disturbing the bed or banks of a stream with a classification and standard of C(t) or higher. For example, 1) the construction of a bridge or placement of a culvert to allow access across a stream; 2) any type of stream bank protection, e.g. placement of rip-rap, or other revetment; 3) lowering stream banks to establish a stream crossing (i.e. creation of a ford); 4) using equipment to remove debris in a stream, all require a permit.

Some examples of activities which are exempt from the requirement to obtain a Protection of Waters permit would be: 1) agricultural activities involving the crossing and recrossing of a stream by livestock or rubber tired farm equipment at an established crossing; or 2) removal of fallen tree limbs or trunks where material can be cabled and pulled from the

stream without disruption of the stream bed or banks, using equipment placed on or above the stream bank. There are occasions when permits from other state or local agencies are required; county or town permits, flood plain permits or other approvals may be necessary. The appropriate offices should be consulted. There is no charge for the Protection of Waters Permit. For permit applications and any questions regarding the permit process contact:

NYSDEC Region 4

Bureau of Habitat

65561 St Hwy 10

Stamford, NY 12167

(607) 652-7741

<http://www.dec.state.ny.us/website/reg4/index.html>

Living Streamside in the East Kill:

Frequently Asked Questions about Working In/Near the Stream

Everyone wants their stream to look and be healthy. Stream health can be measured ecologically by the plants and animals that live in it, but also by its riparian (streamside) buffer area and the stability of its bed and banks. A stable stream is one that does not undergo accelerated erosion. This means the stream does not move laterally (the banks remain stable) or vertically (the stream bed does not build up or cut down) over short periods of time. Streams are very sensitive to anthropogenic (man-made) disturbances, and if stream related projects do not take the necessary precautions, a stable stream can quickly become unstable. Experience has shown that many stream related projects (such as flood control or stream bank stabilization) that have been performed in the past have done far more harm than good to the nation's waterways. Studies that have focused on some of these projects have contributed to the development of new technology to better work with the natural ability of streams to remain stable over time.

Following are answers to some of the questions most commonly asked by homeowners about activities they are considering undertaking that may impact the health and stability of streams. Where you may need more information, contacts are provided. Please contact your local Soil and Water Conservation District office for site-specific information.

We have also noted those activities that may not be beneficial to overall stream health. This information constitutes some of the best professional guidance available today.

If you seek to:

1) Construct a private bridge for vehicles or foot-traffic over the stream, or install a culvert under a driveway or along a stream

Resource Guidance: Efforts should be made to avoid widening or narrowing the stream beyond its naturally stable width. Often, you can observe stable conditions in a reach nearby. Each stream has a stable set of dimensions (width, depth and cross sectional area), which are necessary to maintain effective sediment and water transport. Widening or narrowing can lead to stream instability that could also eventually undermine the bridge. To minimize the potential for erosion or other problems, try to locate a bridge at a narrow and straight reach, and not on a bend. A bridge functions much better than a culvert as a stream crossing, so bridges are preferable to culverts wherever possible. A bridge should span the entire stream to reduce potential erosion damages and prevent debris from catching on the bridge in a flood. If a culvert is absolutely necessary, the size and placement are critical to maintaining stream stability and ensuring the culvert stays in place and minimizes impact on fish passage. DEC's Habitat Unit staff can advise you on size and placement. Multiple culverts (two or more) are rarely permitted.

Permits: Depending on the specific conditions of a stream crossing (bridge or culvert) project, permits are required from the Army Corps of Engineers (ACOE), the New York State Department of Environmental Conservation (DEC) and the New York City Department of Environmental Protection (DEP). An ACOE permit is required when more than 25 cubic yards of fill material will be used below the "ordinary high water mark" (the approximate yearly flood level). Because the streambed or banks will be disturbed, stream crossing construction requires an Article 15 Stream Disturbance Permit from the DEC. Depending upon whether or not there are any drainage features (streams or wetlands) on the property that will be involved as a result of the project, it may require a Crossing, Piping and Diversion Permit (DEP). Also, if the bridge is part of new construction that involves disturbance of more than 1 acre, it must be reviewed under the DEC stormwater State

Pollution Discharge Elimination System (SPDES) program. If the project will disturb more than 2 acres, it may need a Stormwater Pollution Prevention Permit (SPPP) from DEP.

Contacts: Start by contacting the DEC Habitat Unit staff to determine which state permits are needed. In Region 4 (Greene, Schoharie and Delaware Counties), contact Jerry Fraine at 607-652-7366. For DEC Stormwater permits in Region 4 contact Peter Freehafer at 518-357-2381 and at DEP, contact Brenda Drake at 845-340-7633.

2) Divert water from a stream

Resource Guidance: Any diversion of water from a stream, especially during warmer summer months, can negatively impact downstream ecology by reducing the amount of cool water available to aquatic life. This condition can be especially urgent when streamflows are naturally at their lowest levels and trout are in survival-mode. Improper installation of pumps or waterlines can also disturb the streambed or banks, and potentially initiate erosion problems that can worsen over time and move up and downstream to neighboring properties. Finally, water taken from the stream for use nearby will eventually return to the stream, often warmer or containing substances (i.e., lawn chemicals, salts, oils or soap from cars or driveways) that may further stress fish and other aquatic life, or reduce water quality for downstream users.

Permits: Any diversion must be reviewed by DEC.

Contacts: Contact the DEC Habitat Unit. In Region 4 (Greene, Schoharie and Delaware Counties), contact Jerry Fraine at 607-652-7366.

3) Pave or repave a driveway near a stream

Resource Guidance: By not allowing water to slow down and percolate into the ground, impervious surfaces (i.e., pavement and buildings) and associated land drainage improvements that occur from development can accelerate rain runoff into streams, changing the amount and timing of water they receive and in effect delivering it all in one big “gush”. Generally, by the time a watershed exceeds approximately 10% impervious land cover, the streams that capture the runoff are already impaired. A particular concern is localized streambed or bank erosion that a poorly drained impervious surface can encourage. Localized scour and erosion problems can, quickly or slowly, move upstream or downstream

and cause your property or a neighbor's property to erode. Designing "stream friendly" drainage for existing or new impervious surfaces can reduce stream damages from stormwater runoff.

Permits: A DEC Article 15 stream disturbance permit may be required. Seek DEC guidance if the impervious surface is within 50 feet of the stream. If the disturbance is more than 1 acre, it must be reviewed under the DEC stormwater State Pollution Discharge Elimination System (SPDES) program as well. If the project will disturb more than 2 acres, it may need a Stormwater Pollution Prevention Permit (SPPP) from DEP. New driveways being paved for the first time will be required to have a setback from the stream under DEP's regulations.

Contacts: Start by contacting the DEC Habitat Unit to determine what state permits are needed. In Region 4 (Greene, Schoharie and Delaware Counties), contact Jerry Fraine at 607-652-7366. For DEC Stormwater permits, in Region 4, contact Peter Freehafer at 518-357-2381, and at DEP, contact Brenda Drake at 845-340-7633.

4) Cut or trim streamside (riparian) vegetation on the streambank

Resource Guidance: Stable streambanks in the Catskills usually require woody vegetation. Shrub and tree roots provide holding power for streambank soils that cannot be achieved solely by grasses or herbs. For a more thorough discussion on the role of vegetation in stabilizing streambanks, see Section 2.7. To maximize stream bank stability as well as ecological and aesthetic benefits of riparian vegetation, discontinue mowing and allow a buffer of vegetation to grow, or plant woody vegetation.

If you are removing a log jam (a pile of trees that have fallen into the stream and are trapping more trees and stream sediment): this requires technical assistance to ensure that the removal process does not initiate new erosion areas upstream or downstream. These jams can cause considerable property damage. While biologically they may actually be beneficial to the stream, resource management agencies understand the property damage they can cause, and will work with you towards the most beneficial solution. If you are removing individual trees, they must be cut up into smaller pieces and removed from the stream so they will not get caught further downstream and cause or worsen another log or debris jam. If the

log jam or falling trees are not on your property, but are causing damage to your property, you must coordinate with your neighbor.

Permits: The DEC will require an Article 15 Stream Disturbance Permit if the project will disturb the bed or banks of the stream.

Contacts: Seek technical assistance from the DEC Habitat Unit. In Region 4 (Greene, Schoharie and Delaware Counties), contact Jerry Fraine at 607-652-7366. DEP Stream Management Program staff can provide assistance, contact Beth Reichheld at 845-340-7512, or contact your local Soil and Water Conservation District: Greene County SWCD, Rene Van Schaack at 518-622-3620.

5) Stabilize an eroding streambank

Resource Guidance: Streambank stabilization is a common need in the Schoharie valley. As the management plan has revealed, there are eroding streambanks that threaten water quality, private property and public and private infrastructure (i.e., bridges, culverts and roads). Care should be taken in designing stabilization work to ensure that you don't over-widen, narrow, or encroach upon the stream. Borrowing fill material from nearby gravel bars in the stream should be avoided (see FAQ #7). Seek technical assistance to identify the set of causes of your streambank instability problem so the solution can address these causes, and seek a solution that does not transfer the erosion problem up or downstream. The agencies referenced below can advise you on streambank stabilization projects. Neighboring properties may need to be involved to properly address the erosion concern.

Permits: Streambank stabilization will require a DEC Article 15 Stream Disturbance Permit. An ACOE permit is required when more than 25 cubic yards of fill material will be used below the "ordinary high water mark" (the approximate yearly flood level); the DEC can advise you about determining these limits.

Contacts: Seek technical assistance from the DEC Habitat Unit. In Region 4 (Greene, Schoharie and Delaware Counties), contact Jerry Fraine at 607-652-7366. DEP Stream Management Program staff can provide assistance, contact Beth Reichheld at 845-

340-7512, or contact your local Soil and Water Conservation District: Greene County SWCD, Rene Van Schaack at 518-622-3620.

6) Build a house or other structure

Resource Guidance: Siting a new home near a stream can define your enjoyment of that stream and relationship to it. Proper location for homes and facilities must consider stream flooding behavior, no matter how high above or far back from the stream the location may appear during low flows. Because some areas on the FEMA floodplain maps may contain errors due to stream channel migration or infrastructure changes over time, technical assistance is necessary to identify approximate floodplain boundaries, and design your site in as “stream friendly” a manner as possible. Give the stream area to flood, and to move (because a slow rate of erosion is a natural stream adjustment process), so you’ll be able to enjoy living streamside, as well as reducing home maintenance costs from streambank erosion or flood inundation.

Permits: Of course, many permits are needed for new construction, and listing them is beyond the scope of this guidance document. If the house or structure is within 50 ft of a streambank, contact DEC to determine if an Article 15 stream disturbance permit is needed. If the house or driveway will be within 100 ft. of a perennial (flows all year round) stream, you’ll need an Individual Stormwater Permit (DEP). If your project is to construct a single family residence and it will disturb more than 1 acre of land, you must submit a notice of intent to work and an erosion control plan to the DEC under their Stormwater State Pollution Discharge Elimination System (SPDES) program. If your project will disturb more than 2 acres, you’ll need a Stormwater Pollution Prevention Permit (DEP). You will also need to follow State and local regulations, and should contact your Town code enforcement officer. In many communities, the building inspector serves in this capacity.

Contacts: For DEC Article 15 permits: In Region 4 (Greene, Schoharie and Delaware Counties), contact Jerry Fraine at 607-652-7366. For DEC Stormwater permits, in Region 4, contact Peter Freehafer at 518-357-2381, and for DEP permits: Brenda Drake, 845-340-7633. Contact your Town clerk for the number of the local code enforcement officer, and/or building inspector

7) **Extract gravel from the stream**

Resource Guidance: There is a common belief that cleaning gravel from streams is necessary to improve flood conveyance capacity and reduce flooding. Others wish to use skimmed stream gravel for construction-related projects. Proponents of gravel mining should reflect on stream processes including the concept that a stream must effectively be able to move both water and sediment delivered from its watershed to maintain its shape and provide optimum water quality and aquatic habitat. Therefore, any stream channel alterations should consider the impact not only on moving water, but also on sediment (the gravel) transport, to ensure these qualities of a functioning stream are preserved. Excavating gravel usually disturbs the sensitive balance the stream maintains between its slope (steepness) and the amount and size of sediment it can move. Gravel mining reduces the amount of bed material available in the stream system, as a result the stream begins to erode its bed and banks in efforts to bring its sediment load back into balance with its slope and the amount of water in the stream. Gravel mining typically results in accelerated erosion and deposition processes that harm fish habitat. If you are removing gravel to increase flood conveyance capacity, please bear in mind that this has been found to be a damaging practice. If you are excavating gravel for construction-related projects, a non-stream source should be considered.

Permits: DEC rarely permits gravel removal. Any removal will require a DEC Article 15 Stream Disturbance Permit. An ACOE permit is required when more than 25 cubic yards of fill material will be used below the “ordinary high water mark” (the approximate yearly flood level). The DEC can advise you about the need for an ACOE permit.

Contacts: Start by contacting the DEC Habitat Unit to determine what state permits are needed. In Region 4 (Greene, Schoharie and Delaware Counties), contact Jerry Fraine at 607-652-7366. You can also seek technical assistance from the DEP and/or your local Soil and Water Conservation District: Greene County SWCD, Rene Van Schaack at 518-622-3620 and the DEP Stream Management Program, contact Beth Reichheld at 845-340-7512.

Additional Frequently Asked Questions

From: A Guide to Living in Harmony with Streams by the Chemung County SWCD,

<http://www.chemungcountyswcd.com/Tire%20Page.htm>

Who owns the streambed?

New York State is the sovereign owner of the beds of “navigable waters” in the state. This ownership gives the state the right to control the bed and to ensure that navigable waterways shall forever remain public highways. A stream and any contiguous wetlands may be classified as “navigable” if it is large enough for operation of a canoe or larger boat. For information about state ownership of a waterway and the activities for which state approval is required, contact the Lands Underwater program of the NYS Office of General Services (<http://www.ogs.state.ny.us/realEstate/permits/luwfaq.html>). As a general rule, the ownership and therefore control of the bed of non-navigable streams or other non-navigable bodies of water is vested in the proprietors of the adjoining uplands, unless their deed provides otherwise. In other words, if you own the bank of a non-navigable stream, you probably own the streambed and are referred to as a riparian owner. Regardless of who owns a stream, various government entities retain police power over activities that may impact navigation, public safety, the environment, or the rights of other property owners. Owning a stream does not give you the right to do whatever you please with it.

Who owns the water in a stream?

In New York State, water in a stream is not “owned” by anyone. The relevant question is: Who has the right to use water in a stream? Water rights and water laws vary from state to state. New York follows the riparian rights doctrine developed under common law. Common law means that the rules were not enacted by the legislature, but were developed by the courts through the decisions they hand down. Riparian rights doctrine allows the owners of land bordering on a watercourse to withdraw a “reasonable” amount of water. The courts have generally held that domestic use or use on the land is “reasonable,” while removal of water from the riparian property is “unreasonable.” Because all landowners along a stream have “riparian rights,” none can use the water so as to deprive the others of their rights. If a water use interferes with the “reasonable” use of another riparian owner, the aggrieved party must go to court to protect his/her rights.

Who is responsible for the stream?

Restoration of stream problems is generally the responsibility of the private landowner. Although various government agencies have regulatory jurisdiction over how a stream is managed, it is not their job to come and “fix” your stream. Government highway departments generally limit their stream work to that needed for protection of roads, bridges, and culverts. Other government resources are more likely to be available to assist with a project that restores a degraded stream system, rather than one designed for localized protection of private property. For information about stream maintenance and restoration assistance, contact the Greene County Soil and Water Conservation District (518-622-3620). Responsibility for a stream does not give you the right to do whatever you consider necessary to “fix” its problems. Assume that every stream is regulated unless you determine otherwise.

LIABILITY

Common Law is that body of law developed from judicial decisions, based on custom and precedent. As such, it is constantly changing by extension or by interpretation. The central point of common law is damage. The owner of a bridge, hydraulic structure, or other stream project has a legal obligation to protect adjacent landowners from damages due to changes in natural drainage that result from that project. Anyone claiming such damage may file suit in court.

If flooding occurs or gets worse after a stream has been modified (by diverting flow, modifying the channel, constructing a bridge, etc.), is the person who made the modification liable for damages?

Yes, quite possibly. Courts have, according to common law, followed the adage “use your own property in such a manner as not to injure that of another.” This means that no landowner, public or private, has a right to use his/her land in a way that substantially increases flood or erosion damages on adjacent lands. A municipality or property owner may thus be liable for construction, improvements, or modifications that they should reasonably have anticipated to cause property damage to adjacent property. The lack of proper planning, design, and execution thereof, may be considered a clear indication of the lack of good faith and hence negligence with regard to damages that subsequently occurred.

May someone be held liable for failing to remedy a natural hazard that damages adjacent property?

Sometimes. Courts have generally not held governmental units and private individuals responsible for naturally occurring hazards such as stream flooding or bank erosion that damage adjacent lands. In keeping with this principle, a municipality would not be liable for failure to restrain waters between banks of a stream or failure to keep a channel free from obstruction that it did not cause. However, a small number of courts have held that government entities may need to remedy hazards on public lands that threaten adjacent lands. In addition, land owners and governments are liable if they take actions that increase the hazards.

Can liability arise from failure to reasonably operate and maintain a bridge, drainage structure, dam, or flood control structure?

Possibly. The owner of a dam or other water control structure is responsible for inspecting and maintaining it. Where there is a duty to act and the risk of not acting is reasonably perceived, then failure to take appropriate actions may be considered negligent conduct.

May a regulatory agency be liable for issuing a regulatory permit for an activity that damages other private property?

Yes, quite possibly. In fact a careful analysis of hundreds of cases in which the lawsuit involved permitting indicates that a municipality is vastly more likely to be sued for issuing a permit for development that causes harm than for denying a permit based on hazard prevention regulations. The likelihood of a successful lawsuit against a municipality for issuing a permit increases if the permitted activity results in substantial flood, erosion or physical damage to other private property owners.

How safe is safe enough? Municipalities regularly issue permits for activities that are in compliance with existing laws, but might still be at risk of damage.

For example, floodplain development regulations generally apply only to areas mapped as the 100-year floodplain. Yet significant flooding and erosion damages can and do occur outside of these regulated flood-prone areas. Some municipalities address this

additional risk by attaching conditions to their approvals for those projects with identified risks. These conditions can clearly state that the municipality is not obligated to fix personal property in the event of damage. One Town granted approval for a driveway bridge that met all applicable standards, but attached material clearly warning the applicant about the hazards of driving through floodwaters, the risk that emergency vehicles may be unable to reach the house during floods, the potentially high maintenance costs, and the potential liability for the owner if the project results in damage to other property.

May governmental units be held liable for refusing to issue permits in floodways or high-risk erosion areas because the proposed activities could damage other lands?

No. In general, landowners have no right to make a “nuisance” of themselves. Courts have broadly and consistently upheld regulations that prevent one landowner from causing a nuisance or threatening public safety.

What precautions can be taken to avoid liability?

Be “reasonable.” The overall issue, in most instances, is the “reasonableness” of an action by the community or property owner. Due to advances in technology and products, there is an increasingly high standard of care for “reasonable conduct.” The “act of God” defense is seldom successful because even rare flood events are now predictable. As a precaution, technical assistance from stream professionals should be obtained prior to implementing any stream project. Because a well-designed project is less likely to damage other lands, this reduces the potential basis for legal action. And if you are sued, the best defense is a well-documented record showing “due diligence.” That is, that you have done sufficient analysis and design to demonstrate the adequacy of the project with “a reasonable degree of certainty.”